

2016

Mercury in Fish

In Alberta Water Bodies 2009–2013

Alberta
Government



**For more information on Fish Consumption Advisories
Contact:**

Health Protection Branch
Alberta Health
P.O. Box 1360, Station Main
Edmonton, Alberta, T5J 1S6
Telephone: 1-780-427-1470

ISBN: 978-0-7785-8283-0 (Report)
ISBN: 978-0-7785-8284-7 (PDF)

© 2016 Government of Alberta

Executive Summary

Mercury enters the environment through various natural processes and human activities. Methylmercury is transformed from inorganic forms of mercury via methylation by micro-organisms in natural waters, and can accumulate in some fish. Humans are exposed to very low levels of mercury directly from the air, water and food. Fish consumers may be exposed to relatively higher levels of methylmercury by eating mercury-containing fish from local rivers and lakes. Methylmercury can accumulate in the human body over time. Because methylmercury is a known neurotoxin, it is necessary to limit human exposure.

From 2009 to 2013, the Departments of Environment and Parks (AEP) and Health (AH) initiated a survey of mercury levels in fish in selected water bodies in Alberta. These water bodies are extensively accessed by the public for recreational activities.

This report deals with (1) concentrations of total mercury levels in various fish species collected from the water bodies in Alberta, (2) estimated exposures, (3) fish consumption limits, (4) fish consumption advisories, and (5) health benefits of fish consumption. The results indicate that:

1. Concentrations of total mercury in fish in the water bodies in Alberta were within the ranges for the same fish species from other water bodies elsewhere in Canada and the United States.
2. The estimated human exposures to mercury were high for the high fish intake group (over 100 grams/day) who consume walleye, northern pike, lake trout, and lake whitefish from some rivers, lakes and reservoirs in Alberta.
3. Restriction of consumption of walleye, northern pike, lake trout, and lake whitefish from some water bodies was indicated by the health risk assessment, especially for women of reproductive age, pregnant women and young children.
4. Fish consumption advisories are voluntary measures to reduce potential health risk to local fish consumers. The balance between risk and benefits of consumption of mercury-containing fish needs to be understood and considered by consumers.

The Science Advisory Committee reviewed this document and made recommendations. The Public Health Management Committee made the final decisions on fish consumption advisories and measures to inform the public accordingly.

Acknowledgments

Working Group

Health Protection Branch
Alberta Health

Fisheries Management Policy
Alberta Environment and Sustainable Resource and Development

Science Advisory Committee

Dr. Steve E. Hrudey	University of Alberta (Professor Emeritus)
Dr. X. Chris Le	University of Alberta
Dr. Stephan Gabos	University of Alberta
Dr. David Kinniburgh	Alberta Centre for Toxicology

Public Health Management Committee

Office of the Chief Medical Officer of Health
Alberta Health

Health Protection Branch
Alberta Health

Fisheries Management Policy
Alberta Environment and Parks

Environmental Health Team
Alberta Environment and Parks

Senior Medical Officer of Health
Alberta Health Services

First Nations and Inuit Health Branch
Health Canada

Table of Contents

1. Introduction	1
2. Materials and Methods	3
2.1 Units Used for Expressing Mercury Data	3
2.2 Field Collection	3
2.3 Laboratory Analysis	14
2.4 Estimation of Exposure Ratio	14
2.5 Consumption Limits	15
3. Results and Discussions	16
3.1 Concentrations in Fish	16
3.2 Local Fish Consumption Rates	29
3.3 Estimated Exposures	30
3.4 Consumption Limits	41
3.5 Fish Consumption Advisories	44
3.6 Benefits of Fish Consumption	45
3.7 Criteria for Issuing Fish Consumption Advisories	47
4. Conclusions	48
5. References	52
Appendix A Data from Environment Canada	59
Appendix B Guide for Use of Food Consumption Advisory Information	63

List of Tables

Table 1 Units Used for Expressing Mercury Data related to Fish	3
Table 2 Sample Size and Mean of Weight and Length, 2009/2010	9
Table 3 Sample Size and Mean of Weight and Length, 2010/2011	10
Table 4 Sample Size and Mean of Weight and Length, 2011/2012	11
Table 5 Sample Size and Mean of Weight and Length, 2012/2013	12
Table 6 Total Mercury Levels in Fish, 2009/2010 (µg/g, wet weight)	17
Table 7 Total Mercury Levels in Fish, 2010/2011 (µg/g, wet weight)	19
Table 8 Total Mercury Levels in Fish, 2011/2012 (µg/g, wet weight)	21
Table 9 Total Mercury Levels in Fish, 2012/2013 (µg/g, wet weight)	23
Table 10 Mean THg Concentrations in Fish Muscles in Alberta Water Bodies ...	27
Table 11 Mean THg Levels in Fish Muscles Reported in the Literature	28
Table 12 Local Fish Consumption Rates in Communities of Central Alberta	29
Table 13 Estimated Exposure Ratios for Women of Reproductive Age ^a	32
Table 14 Estimated Exposure Ratios for Adults	37
Table 15 Lifetime Fish Consumption Limits ^a	42
Table 16 Recommended Fish Consumption Limits	49

List of Figures

Figure 1 Sampling Locations for 2009/2010 and 2010/2011	7
Figure 2 Sampling Locations for 2011/2012 and 2012/2013	8
Figure 3 Mean Concentrations of Total Mercury in Fish, 2009/2010	18
Figure 4 Mean Concentrations of Total Mercury in Fish, 2010/2011	20
Figure 5 Mean Concentrations of Total Mercury in Fish, 2011/2012	22
Figure 6 Mean Concentrations of Total Mercury in Fish, 2012/2013	24

1. Introduction

Mercury (Hg) occurs naturally in the environment. There are three forms of mercury: elemental (metallic) mercury, inorganic mercury salts and organic mercury compounds. Mercury enters the environment through natural processes and human activities. The form of mercury most commonly found in the air is volatile elemental mercury. Methylmercury (MeHg) is often formed from other forms of mercury during natural biological processes such as methylation by micro-organisms in the water and sediment. MeHg can accumulate in some fish. People are exposed to very low levels of mercury in the air, water and food. Some people may be exposed to relatively higher levels of MeHg through eating mercury-containing fish. MeHg accumulates in the human body over time. Because MeHg is a known neurotoxin, it is necessary to limit human exposure.

To protect public health, Health Canada has proposed mercury guidelines, and advisories for different fish consumer groups (Health Canada, 1979; Feeley and Lo, 1998; Health Canada 2007; Feeley 2008) based on total mercury (THg) or MeHg. These values are expressed either in units of μg THg or MeHg per g of fish flesh or as a Provisional Tolerable Daily Intake (pTDI) in units of μg THg or MeHg per kg of consumer body mass per day (see Section 2.1).

1. $0.5 \mu\text{g}$ THg/g for all commercial fish/seafood (Guideline);
2. $1.0 \mu\text{g}$ THg/g for certain commercial fish species such as fresh and frozen tuna, shark, swordfish, escolar, marlin and orange roughy which are known to be consumed less frequently (Advisory); and
3. $0.2 \mu\text{g}$ THg/g for subsistence consumers (Advisory)
4. $0.2 \mu\text{g}$ MeHg/kg bwt pTDI for women of reproductive age and young children (Guideline);
5. $0.47 \mu\text{g}$ MeHg/kg bwt pTDI for the general population (Guideline)

The guidelines for commercial fish/seafood are used as a general screening criterion, with the knowledge that most species of commercial fish usually contain lower levels ($< 0.1 \mu\text{g/g}$) of mercury. This guideline is enforceable by the Canadian Food Inspection Agency (CFIA) for commercial fish. For example, the CFIA has been monitoring total mercury (THg) levels in commercial fish caught from Lake Athabasca in Alberta since the early 1990s. The recommendation for subsistence consumers proposed by Health Canada since the 70s is used for First Nations and Inuit people relying on subsistence fresh water fishing when Medical Services Branch became aware of long term fish consumption patterns of over 100 g/d (Health Canada 1979). The First Nations and Inuit subsistence consumers should limit their fish consumption if the mercury levels in fish are over $0.2 \mu\text{g}$ THg/g. Over $0.5 \mu\text{g}$ THg/g fish consumption should be avoided for any consumer.

Fish consumption advisories are developed based on these pTDIs. These advisories provide the public with a warning of potential health risk resulting from consuming

total mercury-containing fish. Fish consumption advisories are designed to minimize the potential health risks to fish consumers who eat voluntarily selected fish consumption.

From 2005 to 2013, the Departments of Environment and Parks (AEP) and Health (AH) initiated a survey of mercury levels in fish in selected water bodies in Alberta. These water bodies are extensively accessible to the public for recreational activities.

Environment Canada sampled and tested fish from Alberta lakes for mercury during the same time period. The results of these analyses are included in this report.

The results related to mercury in fish are discussed as follows:

1. mercury concentrations in fish;
2. comparison of mercury concentrations in the same fish species in the rivers and lakes in Canada and the U.S.;
3. total fish consumption rates;
4. estimated exposures for women of reproductive age, children and adults;
5. fish consumption advisories; and
6. health benefits of fish consumption.

2. Materials and Methods

2.1 Units Used for Expressing Mercury Data

A summary of the different units that may be used for expressing mercury concentrations is presented in Table 1. For the purposes of this report, the following conversion of values reported from different sources, all data on mercury concentration is first put in equivalent mg kg of mercury per g of fish, i.e. mg kg⁻¹ (equivalent to mg kg⁻¹ of mercury per million parts of fish (ppm)). Likewise, fish mercury exposure will be expressed as kg of mercury per kg of fish per body mass per day, i.e. kg kg⁻¹ day⁻¹. Comparisons between all the information from fish mercury exposure levels will be expressed as g of fish consumed per week, i.e. g wk⁻¹.

Table 1
Units Used for Expressing Mercury Data, Abbreviated as mg.

Parameter	Traditional Unit	Alternative Unit	Equivalent Unit
mg Concentration	kg of fish per g of fish wet weight ppm	kg of fish per kg of fish wet weight mg/kg	1 mg/kg wet weight ppm (1000 ppb)
mg/kg mercury/kg fisherman	kg of fish per kg of fisherman body weight (ppm) per kg of fish per kg fisherman		
Recommended for comparisons (ppm)	g of mercury containing 100 mg fisherman per week g/100	g of mercury containing 100 mg consumed per week g/100	1 g/100 g fish

2.2 Field Collection

Field collection of 10 sampling sites (Figure 1) was conducted by VCC personnel September 2009 and January 2010. Sampling sites included:

1. Caring Lake
2. Chagaming Lake (South side area)
3. Cowan Reservoir
4. Crossway Valley Reservoir
5. Egan Lake
6. Hay River at mouth of Chagaming River
7. Koo Lake
8. McLaughlin Lake
9. Lake Athabasca
10. Lake St. Charles Reservoir
11. Rutherford Lake
12. Stirling Hills Reservoir

- 10. Muskegon Lake
- 11. Torch Lake
- 12. Torch Lake
- 13. Torch Lake

These collections are being reviewed, and Figure 2 will be updated, with additional
collections, 2000 and 2001. Sampling sites included:

- 1. Torch Lake
- 2. Torch Lake
- 3. Torch Lake
- 4. Torch Lake
- 5. Torch Lake
- 6. Torch Lake
- 7. Torch Lake
- 8. Torch Lake
- 9. Torch Lake
- 10. Torch Lake
- 11. Torch Lake
- 12. Torch Lake
- 13. Torch Lake
- 14. Torch Lake
- 15. Torch Lake
- 16. Torch Lake
- 17. Torch Lake
- 18. Torch Lake
- 19. Torch Lake
- 20. Torch Lake
- 21. Torch Lake
- 22. Torch Lake
- 23. Torch Lake
- 24. Torch Lake
- 25. Torch Lake
- 26. Torch Lake
- 27. Torch Lake
- 28. Torch Lake
- 29. Torch Lake
- 30. Torch Lake
- 31. Torch Lake
- 32. Torch Lake
- 33. Torch Lake
- 34. Torch Lake
- 35. Torch Lake
- 36. Torch Lake
- 37. Torch Lake
- 38. Torch Lake
- 39. Torch Lake
- 40. Torch Lake
- 41. Torch Lake
- 42. Torch Lake
- 43. Torch Lake
- 44. Torch Lake
- 45. Torch Lake
- 46. Torch Lake
- 47. Torch Lake
- 48. Torch Lake
- 49. Torch Lake
- 50. Torch Lake
- 51. Torch Lake
- 52. Torch Lake
- 53. Torch Lake
- 54. Torch Lake
- 55. Torch Lake
- 56. Torch Lake
- 57. Torch Lake
- 58. Torch Lake
- 59. Torch Lake
- 60. Torch Lake
- 61. Torch Lake
- 62. Torch Lake
- 63. Torch Lake
- 64. Torch Lake
- 65. Torch Lake
- 66. Torch Lake
- 67. Torch Lake
- 68. Torch Lake
- 69. Torch Lake
- 70. Torch Lake
- 71. Torch Lake
- 72. Torch Lake
- 73. Torch Lake
- 74. Torch Lake
- 75. Torch Lake
- 76. Torch Lake
- 77. Torch Lake
- 78. Torch Lake
- 79. Torch Lake
- 80. Torch Lake
- 81. Torch Lake
- 82. Torch Lake
- 83. Torch Lake
- 84. Torch Lake
- 85. Torch Lake
- 86. Torch Lake
- 87. Torch Lake
- 88. Torch Lake
- 89. Torch Lake
- 90. Torch Lake
- 91. Torch Lake
- 92. Torch Lake
- 93. Torch Lake
- 94. Torch Lake
- 95. Torch Lake
- 96. Torch Lake
- 97. Torch Lake
- 98. Torch Lake
- 99. Torch Lake
- 100. Torch Lake

These collections are being reviewed, and Figure 2 will be updated, with additional
collections, 2000 and 2001. Sampling sites included:

- 1. Torch Lake
- 2. Torch Lake
- 3. Torch Lake
- 4. Torch Lake
- 5. Torch Lake
- 6. Torch Lake
- 7. Torch Lake
- 8. Torch Lake
- 9. Torch Lake
- 10. Torch Lake
- 11. Torch Lake
- 12. Torch Lake
- 13. Torch Lake
- 14. Torch Lake
- 15. Torch Lake
- 16. Torch Lake
- 17. Torch Lake
- 18. Torch Lake
- 19. Torch Lake
- 20. Torch Lake
- 21. Torch Lake
- 22. Torch Lake
- 23. Torch Lake
- 24. Torch Lake
- 25. Torch Lake
- 26. Torch Lake
- 27. Torch Lake
- 28. Torch Lake
- 29. Torch Lake
- 30. Torch Lake
- 31. Torch Lake
- 32. Torch Lake
- 33. Torch Lake
- 34. Torch Lake
- 35. Torch Lake
- 36. Torch Lake
- 37. Torch Lake
- 38. Torch Lake
- 39. Torch Lake
- 40. Torch Lake
- 41. Torch Lake
- 42. Torch Lake
- 43. Torch Lake
- 44. Torch Lake
- 45. Torch Lake
- 46. Torch Lake
- 47. Torch Lake
- 48. Torch Lake
- 49. Torch Lake
- 50. Torch Lake
- 51. Torch Lake
- 52. Torch Lake
- 53. Torch Lake
- 54. Torch Lake
- 55. Torch Lake
- 56. Torch Lake
- 57. Torch Lake
- 58. Torch Lake
- 59. Torch Lake
- 60. Torch Lake
- 61. Torch Lake
- 62. Torch Lake
- 63. Torch Lake
- 64. Torch Lake
- 65. Torch Lake
- 66. Torch Lake
- 67. Torch Lake
- 68. Torch Lake
- 69. Torch Lake
- 70. Torch Lake
- 71. Torch Lake
- 72. Torch Lake
- 73. Torch Lake
- 74. Torch Lake
- 75. Torch Lake
- 76. Torch Lake
- 77. Torch Lake
- 78. Torch Lake
- 79. Torch Lake
- 80. Torch Lake
- 81. Torch Lake
- 82. Torch Lake
- 83. Torch Lake
- 84. Torch Lake
- 85. Torch Lake
- 86. Torch Lake
- 87. Torch Lake
- 88. Torch Lake
- 89. Torch Lake
- 90. Torch Lake
- 91. Torch Lake
- 92. Torch Lake
- 93. Torch Lake
- 94. Torch Lake
- 95. Torch Lake
- 96. Torch Lake
- 97. Torch Lake
- 98. Torch Lake
- 99. Torch Lake
- 100. Torch Lake

These collections are being reviewed, and Figure 2 will be updated, with additional
collections, 2000 and 2001. Sampling sites included:

These collections are being reviewed, and Figure 2 will be updated, with additional
collections, 2000 and 2001. Sampling sites included:

© 2000 Blackwell Science Ltd, *Journal of Internal Medicine* 247: 399–405

© 2000 Blackwell Science Ltd, *Journal of Internal Medicine* 247: 395–402

This journal has been peer-reviewed by the following referees:
Dr. J. H. M. van der Kolk



Figure 1 Sampling Locations for 2010/2011 and 2011/2012



Figure 2 Sampling Locations for 2011/2012 and 2012/2013

Table 2 Sample Size and Mean of Weight and Length, 2009/2010

Lake and Species	Sample Size	Length (cm)	Wet Weight (g)
<u>Callina Lake</u>			
Northern pike	7	67	1,964
Walleye	10	54	1,456
<u>Christina Lake</u>			
Brook trout	15	33	340
<u>Cowoki Reservoir</u>			
Northern pike	6	75	3,409
Walleye	4	59	2,563
<u>Crimling Valley Reservoir</u>			
Northern pike	8	63	1,588
Walleye	11	48	1,057
<u>Edith Lake</u>			
Brook trout	20	31	352
<u>Hay River</u>			
Walleye	4	49	1,277
<u>Isle Lake</u>			
Northern pike	20	53	1,104
Walleye	21	50	1,247
<u>Kehewin Lake</u>			
Northern pike	10	64	1,635
Walleye	10	45	957
<u>Lake Athabasca</u>			
Lake trout	30	70	2,872
Lake whitefish	4	44	969
Northern pike	29	70	2,936
Walleye	30	50	1,453
<u>Little Bow Reservoir</u>			
Northern pike	9	63	1,489
<u>McGregor Lake</u>			
Northern pike	2	82	4,200
Walleye	12	58	1,893
<u>Rolling Hills Reservoir</u>			
Northern pike	6	80	3,700
Walleye	7	61	2,523
<u>Sturgeon Lake</u>			
Lake whitefish	10	49	1,255
Northern pike	10	61	1,573
Walleye	10	46	917
<u>Touchwood Lake</u>			
Lake whitefish	9	57	1,648
Northern pike	10	86	4,317
Walleye	10	63	2,051
<u>Wabamun Lake</u>			
Lake whitefish	13	47	1,017
Northern pike	10	74	3,092
<u>Whitefish Lake</u>			
Lake whitefish	6	67	3,486
Northern pike	7	81	3,816
Walleye	6	58	1,935

Table 3 Sample Size and Mean of Weight and Length, 2010/2011

<u>Lake and Species</u>	<u>Sample Size</u>	<u>Length (cm)</u>	<u>Wet Weight (g)</u>
<u>Amisk Lake</u>			
Northern pike	11	53	835
Walleye	10	43	665
<u>Bitscho Lake</u>			
Northern pike	10	75	3,226
Walleye	10	46	1,548
<u>Chinchaga River</u>			
Walleye	5	42	718
<u>Dore Lake^a</u>			
Northern pike	20	69.8	2,808
Walleye	20	46.1	1,051
<u>Isle Lake</u>			
Northern pike	10	75	3,092
Lake whitefish	13	47	1,017
<u>Kinnaird Lake</u>			
Northern pike	11	56	1,380
Walleye	10	48	1,080
<u>Lac Bellevue</u>			
Walleye	12	44	912
<u>Laurier Lake</u>			
Northern pike	4	81	3,128
<u>Lein Thompson</u>			
Rainbow trout	10	33	493
<u>Lesser Slave Lake East Basin</u>			
Northern pike	9	69	2,301
Walleye	14	52	1,136
<u>Lesser Slave Lake West Basin</u>			
Northern pike	16	76	2,913
Walleye	15	59	1,155
<u>Long Lake</u>			
Northern pike	6	51	826
Walleye	2	46	953
<u>Meander River</u>			
Northern pike	7	59	1,414
Walleye	3	39	587
<u>North Wabasca Lake</u>			
Northern pike	10	82	4,294
Walleye	10	57	1,746
<u>Pinehurst Lake</u>			
Northern pike	10	55	1,575
Walleye	8	54	1,839
<u>Pitchimi Lake</u>			
Lake trout	15	79	4,671
<u>Richardson Lake</u>			
Northern pike ^a	19	76.0	4,066
Walleye ^a	20	47.4	1,289
Lake whitefish	20	46	1,172
<u>Skeleton Lake</u>			
Northern pike	7	52	845
Walleye	9	51	1,462
<u>Snipe Lake</u>			
Northern pike	10	75	2,969
Walleye	10	55	1,846

Lake and Species	Sample Size	Length (cm)	Wet Weight (g)
<u>Winigami Lake</u>			
Northern pike	11	64	1,533
Walleye	10	59	2,069

^a Data source – Environment Canada

Table 4 Sample Size and Mean of Weight and Length, 2011/2012

Lake and Species	Sample Size	Length (cm)	Wet Weight (g)
<u>Baptiste Lake</u>			
Northern pike	10	67	1,984
Walleye	8	43	738
<u>Beaver Lake</u>			
Lake whitefish	10	52	1,438
Northern pike	11	67	2,210
Walleye	15	51	1,304
<u>Bourque Lake</u>			
Lake whitefish	10	N/A	849
Northern pike	8	N/A	1,502
Walleye	10	N/A	1,701
<u>Calling Lake</u>			
Northern pike	10	64	1,390
Walleye	12	51	1,076
<u>Cross (Steele) Lake</u>			
Northern pike	10	66	1,860
<u>Heart Lake</u>			
Lake whitefish	7	59	2,371
Northern pike	14	73	2,662
Walleye	27	55	1,964
<u>Hutch Lake</u>			
Walleye	10	53	1,993
<u>Kirby Lake</u>			
Lake whitefish	21	55	1,625
Northern pike	20	73	2,784
<u>Loon River</u>			
Walleye	4	55	989
<u>McMillan Lake</u>			
Northern pike	13	70	1,948
<u>Moose Lake</u>			
Northern pike	17	N/A	1,167
Walleye	10	N/A	1,633
<u>Nipisi Lake</u>			
Northern pike	20	74	2,706
<u>Peerless Lake</u>			
Lake trout	9	71	3,824
Walleye	16	55	1,526
<u>Wolf Lake</u>			
Lake whitefish	10	N/A	1,723
Northern pike	10	N/A	1,258
Walleye	10	N/A	973

Table 5 Sample Size and Mean of Weight and Length, 2012/2013

Lake and Species	Sample Size	Length (cm)	Wet Weight (g)
<u>Baptiste Lake</u>			
Lake whitefish	5	40	908
Northern pike	10	60	1,370
Walleye	33	49	1,061
<u>Eagle Lake</u>			
Northern pike	13	49	777
Walleye	23	48	1,323
<u>Elinor Lake</u>			
Lake whitefish	14	51	1,347
Northern pike	8	73	1,625
Walleye	17	53	1,968
<u>Ethel Lake</u>			
Lake whitefish	12	59	2,360
Northern pike	9	54	740
Walleye	19	48	1,009
<u>Goodfish Lake</u>			
Northern pike	7	63	1,541
Walleye	9	50	1,261
<u>Gregoire Lake</u>			
Lake whitefish	4	53	2,243
Walleye	9	50	1,273
<u>Hilda Lake</u>			
Northern pike	8	56	1,128
Walleye		19	46
<u>Keho Lake</u>			
Lake whitefish	30	45	874
Northern pike	30	63	1,506
Walleye			24
<u>Lac la Nonne</u>			
Lake whitefish	15	52	1,634
Northern pike	8	55	1,028
Walleye	17	41	684
<u>Lac Ste Anne</u>			
Walleye	7	43	798
<u>Lake Isle</u>			
Northern pike	12	49	813
<u>Marie Lake</u>			
Lake whitefish	8	45	794
Northern pike	7	66	1,986
Walleye	12	49	1,133
<u>McLeod Lake</u>			
Rainbow Trout	7	31	339
<u>Moonshine Lake</u>			
Rainbow Trout	24	29	294
<u>Moose Lake</u>			
Cisco	26	40	900
Lake whitefish	5	60	2,582
Northern pike	50	66	1,759
Walleye	48	55	1,625
Yellow Perch	36	22	155

Lake and Species	Sample Size	Length (cm)	Wet Weight (g)
<u>Pigeon Lake</u>			
Lake whitefish	38	54	1,934
Walleye	31	49	1,115
<u>Pine Lake</u>			
Northern pike	5	58	1,418
Walleye	5	48	1,112
<u>Rock Island Lake</u>			
Northern pike	17	58	1,290
Walleye	19	51	1,509
<u>Sylvan Lake</u>			
Lake whitefish	24	39	597
Walleye	15	37	584
<u>Whitefish Lake</u>			
Lake whitefish	4	47	962
Northern pike	4	66	1,689
Walleye	16	55	1,526
<u>Wizard Lake</u>			
Northern pike	15	50	841

2.3 Laboratory Analysis

In 2009/2010, laboratory analysis was performed in the Biogeochemical Analytical Laboratory, Department of Biological Sciences at the University of Alberta. In 2010/2011, 2011/2012 and 2012/2013, laboratory analyses were performed by the Alberta Centre for Toxicology at the University of Calgary, Alberta. The analytical method was based on the modified USEPA methods 200.7 and 200.8 (USEPA 2001). The Environment Canada laboratory used the USEPA method 7473 (USEPA 2007) (Appendix A).

2.4 Estimation of Exposure Ratio

Estimated daily intake (EDI), $\mu\text{g/d}$, was calculated as follows:

$$\text{EDI} = C \times \text{IR} \times \text{BF}/\text{BW}$$

C is a representative measure of THg concentrations in fish muscle ($\mu\text{g/g}$). From a human health perspective, the amount of MeHg is of most interest. In mercury analyses of fish, the THg in the sample is measured, which is the sum of all forms of mercury present, rather than MeHg because the analysis of MeHg is more expensive. Some studies reported that the percentage of MeHg in THg in fish ranged from 81% to 95% (CFIA 2003). For the purposes of health risk assessments, 100% of THg is assumed to be MeHg, thereby erring on the side of caution.

IR is the human rate of fish consumption (g/d).

BF is bioavailability factor (assumed to be 100%, unless, the maximum possible and most cautious assumption for this factor).

BW is average body weight in humans (kg). The average of body weight for male and female adults in Alberta is 73 kg. The average human body weights used by Health Canada are 65 kg for women of reproductive age, 26.4 kg for the five to 11 year old group, and 14.4 kg for the one to four year old group (Health Canada 2007).

Exposure ratio (ER, unitless) was calculated by using the following equation:

$$\text{ER} = \text{EDI} / \text{pTDI}$$

The provisional tolerable daily intake (pTDI, $\mu\text{g MeHg/kg bwt/d}$) is determined by toxicological risk assessment on mercury (Health Canada 2007). The pTDI for MeHg is the maximum amount of mercury that can be ingested on a daily basis over a lifetime without increased risk of adverse health effects. Health Canada proposed a pTDI of mercury as 0.2 $\mu\text{g Hg/kg bwt/d}$ for women of reproductive (childbearing) age and for children. Children refer to two age groups: the five to 11 year old group and the one to four year old group. Health Canada proposed a pTDI of MeHg as 0.47 μg

Hg/kg wet wt for adults (adult men and adult women who are not of reproductive age)

2.5 Consumption Limits

For quantitative fish advisories, the lifetime average consumption limits (weekly basis) are calculated. The calculation of the consumption limits (CL, g fish per week) is based on the following equation:

$$CL = pTDI \times BW(7 \text{ d/week}) / C$$

Where pTDI is provisional tolerable daily intake ($\mu\text{g MeHg/kg body wt}$),
BW is body weight (mass) in humans (kg), and
C ($\mu\text{g Hg} / \text{g fish}$) is the measured THg concentration in fish muscle.

The consumption limits that correspond to the Health Canada pTDI and the commercial fish Hg recommendation (maximum concentration of $0.5 \mu\text{g Hg/g fish}$) are provided below as a reference point.

Consumption Limits for adult men and adult women not of reproductive age

$$CL = (0.47 \mu\text{g MeHg/kg body wt})(73 \text{ kg})(7 \text{ d/week}) / (0.5 \mu\text{g Hg} / \text{g fish}) = 480 \text{ g fish/week}$$

Consumption Limits for women of reproductive age

$$CL = (0.2 \mu\text{g MeHg/kg body wt})(65 \text{ kg})(7 \text{ d/week}) / (0.5 \mu\text{g Hg} / \text{g fish}) = 180 \text{ g fish/week}$$

Consumption Limits for children age 5 – 11 (body weight 25.4 kg)

$$CL = (0.2 \mu\text{g MeHg/kg body wt})(25.4 \text{ kg})(7 \text{ d/week}) / (0.5 \mu\text{g Hg} / \text{g fish}) = 74 \text{ g fish/week}$$

Consumption Limits for children age 1 – 4 (body weight 14.4 kg)

$$CL = (0.2 \mu\text{g MeHg/kg body wt})(14.4 \text{ kg})(7 \text{ d/week}) / (0.5 \mu\text{g Hg} / \text{g fish}) = 40 \text{ g fish/week}$$

3. Results and Discussions

3.1 Concentrations in Fish

The total mercury concentrations in wet weight in fish are summarized in Tables 6, 7, 8, and 9 and Figures 3, 4, 5, and 6. The concentrations exceeding the 0.5 $\mu\text{g/g}$ commercial fish limit are shown in **bold**.

The average THg concentrations exceeded 0.5 $\mu\text{g/g}$ (ppm) in northern pike in the following water bodies:

1. Annapolis Lake
2. Cocks Bay Reservoir
3. Elmer Lake
4. Halls Lake
5. McGowan Lake
6. Pinesburg Lake
7. Rolling Hills Reservoir
8. Tappanville Lake
9. Wheelock Lake

The average THg concentrations exceeded 0.5 $\mu\text{g/g}$ (ppm) in catfish in the following water bodies:

1. Annapolis Lake
2. Bayview Lake
3. Cocks Bay Reservoir
4. Crawling Valley Reservoir
5. Dora Lake
6. Elmer Lake
7. Hay River at mouth of Chesapeake River
8. Halls Lake
9. Kinnard Lake
10. Lee in Stone
11. McGowan Lake
12. Pinesburg Lake
13. Rolling Hills Reservoir
14. Tappanville Lake and
15. Wheelock Lake
16. Wolf Lake

The average THg concentrations exceeded 0.5 $\mu\text{g/g}$ (ppm) in lake whitefish from Wheelock Lake and in lake trout in Pinesburg Lake.

Table 6: Total Memory Levels in RAM, 2008-2009

State and District	2008	2009	2010
Alabama			
Alabama	5.15	5.15	5.15
Montgomery	5.15	5.15	5.15
Alaska			
Alaska	5.15	5.15	5.15
Arizona			
Arizona	5.15	5.15	5.15
Phoenix	5.15	5.15	5.15
Arkansas			
Arkansas	5.15	5.15	5.15
Little Rock	5.15	5.15	5.15
California			
California	5.15	5.15	5.15
San Francisco	5.15	5.15	5.15
Colorado			
Colorado	5.15	5.15	5.15
Denver	5.15	5.15	5.15
Connecticut			
Connecticut	5.15	5.15	5.15
Hartford	5.15	5.15	5.15
Delaware			
Delaware	5.15	5.15	5.15
Dover	5.15	5.15	5.15
District of Columbia			
District of Columbia	5.15	5.15	5.15
Florida			
Florida	5.15	5.15	5.15
Tallahassee	5.15	5.15	5.15
Georgia			
Georgia	5.15	5.15	5.15
Atlanta	5.15	5.15	5.15
Hawaii			
Hawaii	5.15	5.15	5.15
Idaho			
Idaho	5.15	5.15	5.15
Illinois			
Illinois	5.15	5.15	5.15
Chicago	5.15	5.15	5.15
Indiana			
Indiana	5.15	5.15	5.15
Indianapolis	5.15	5.15	5.15
Iowa			
Iowa	5.15	5.15	5.15
Des Moines	5.15	5.15	5.15
Kansas			
Kansas	5.15	5.15	5.15
Topeka	5.15	5.15	5.15
Kentucky			
Kentucky	5.15	5.15	5.15
Frankfort	5.15	5.15	5.15
Louisiana			
Louisiana	5.15	5.15	5.15
Baton Rouge	5.15	5.15	5.15
Maine			
Maine	5.15	5.15	5.15
Portland	5.15	5.15	5.15
Maryland			
Maryland	5.15	5.15	5.15
Baltimore	5.15	5.15	5.15
Massachusetts			
Massachusetts	5.15	5.15	5.15
Boston	5.15	5.15	5.15
Michigan			
Michigan	5.15	5.15	5.15
Lansing	5.15	5.15	5.15
Minnesota			
Minnesota	5.15	5.15	5.15
St. Paul	5.15	5.15	5.15
Mississippi			
Mississippi	5.15	5.15	5.15
Jackson	5.15	5.15	5.15
Missouri			
Missouri	5.15	5.15	5.15
Kansas City	5.15	5.15	5.15
Montana			
Montana	5.15	5.15	5.15
Helena	5.15	5.15	5.15
Nebraska			
Nebraska	5.15	5.15	5.15
Lincoln	5.15	5.15	5.15
Nevada			
Nevada	5.15	5.15	5.15
Las Vegas	5.15	5.15	5.15
New Hampshire			
New Hampshire	5.15	5.15	5.15
Manchester	5.15	5.15	5.15
New Jersey			
New Jersey	5.15	5.15	5.15
Newark	5.15	5.15	5.15
New Mexico			
New Mexico	5.15	5.15	5.15
Santa Fe	5.15	5.15	5.15
New York			
New York	5.15	5.15	5.15
New York City	5.15	5.15	5.15
North Carolina			
North Carolina	5.15	5.15	5.15
Raleigh	5.15	5.15	5.15
North Dakota			
North Dakota	5.15	5.15	5.15
Bismarck	5.15	5.15	5.15
Ohio			
Ohio	5.15	5.15	5.15
Columbus	5.15	5.15	5.15
Oklahoma			
Oklahoma	5.15	5.15	5.15
Oklahoma City	5.15	5.15	5.15
Oregon			
Oregon	5.15	5.15	5.15
Portland	5.15	5.15	5.15
Pennsylvania			
Pennsylvania	5.15	5.15	5.15
Philadelphia	5.15	5.15	5.15
Rhode Island			
Rhode Island	5.15	5.15	5.15
Providence	5.15	5.15	5.15
South Carolina			
South Carolina	5.15	5.15	5.15
Columbia	5.15	5.15	5.15
South Dakota			
South Dakota	5.15	5.15	5.15
Sioux Falls	5.15	5.15	5.15
Tennessee			
Tennessee	5.15	5.15	5.15
Nashville	5.15	5.15	5.15
Texas			
Texas	5.15	5.15	5.15
Houston	5.15	5.15	5.15
Utah			
Utah	5.15	5.15	5.15
Salt Lake City	5.15	5.15	5.15
Vermont			
Vermont	5.15	5.15	5.15
Montpelier	5.15	5.15	5.15
Virginia			
Virginia	5.15	5.15	5.15
Richmond	5.15	5.15	5.15
Washington			
Washington	5.15	5.15	5.15
Seattle	5.15	5.15	5.15
West Virginia			
West Virginia	5.15	5.15	5.15
Martinsburg	5.15	5.15	5.15
Wisconsin			
Wisconsin	5.15	5.15	5.15
Milwaukee	5.15	5.15	5.15
Wyoming			
Wyoming	5.15	5.15	5.15
Cheyenne	5.15	5.15	5.15




Figure 1: Percentage of First Arrests by Age Group and Gender

Table 1. Total number of students in each grade, by sex, and by race/ethnicity, 1990-1991

Grade	Male	Female	Total
Kindergarten	1,100	1,100	2,200
Grade 1	1,100	1,100	2,200
Grade 2	1,100	1,100	2,200
Grade 3	1,100	1,100	2,200
Grade 4	1,100	1,100	2,200
Grade 5	1,100	1,100	2,200
Grade 6	1,100	1,100	2,200
Grade 7	1,100	1,100	2,200
Grade 8	1,100	1,100	2,200
Grade 9	1,100	1,100	2,200
Grade 10	1,100	1,100	2,200
Grade 11	1,100	1,100	2,200
Grade 12	1,100	1,100	2,200
Total	13,200	13,200	26,400

Variable	Mean	SD	Range
Age	32.1	7.2	18-55
Relationship duration	5.8	3.5	1-15
Relationship type			
Married	65.2		
Common-law	34.8		



Figure 1: Relationship Characteristics of Participants in the Study (N = 1,000)

Table 6 Total Mercury Levels in Fish, 2010/2012 (ug/g wet weight)

Lake and Species	2010	2011	2012
Barrie Lake			
Northern pike	0.38	0.16	0.72
Walleye	0.21	0.11	0.33
Beaver Lake			
Lake whitefish	0.13	0.08	0.15
Northern pike	0.45	0.21	1.11
Walleye	0.35	0.13	0.49
Brantford Lake			
Lake whitefish	0.37	0.32	0.19
Northern pike	0.33	0.12	0.51
Walleye	0.38	0.28	0.36
Canby Lake			
Northern pike	0.15	0.18	0.49
Walleye	0.13	0.08	0.32
Crook (Shedden) Lake			
Northern pike	0.21	0.07	0.35
Georgian Lake			
Lake whitefish	0.32	0.31	0.36
Northern pike	0.18	0.08	0.23
Walleye	0.11	0.06	0.32
Georgian Lake			
Walleye	0.11	0.08	0.13
Georgian Lake			
Lake whitefish	0.32	0.31	0.05
Northern pike	0.13	0.02	0.36
Georgian Lake			
Walleye	0.38	0.18	0.65
Georgian Lake			
Northern pike	0.35	0.21	0.65
Georgian Lake			
Eastern Trout	0.37	0.06	0.12
Georgian Lake			
Northern pike	0.33	0.05	0.66
Walleye	0.45	0.28	0.67
Georgian Lake			
Northern pike	0.11	0.05	0.18
Georgian Lake			
Lake trout	0.13	0.05	0.21
Georgian Lake			
Lake whitefish	0.13	0.05	0.26
Northern pike	0.39	0.12	0.55
Walleye	0.48	0.21	0.73




Figure 5 Mean Concentrations of Total Mercury in Fish, 2011/2012

Table 9 Total Mercury Levels in Fish, 2012/2013 ($\mu\text{g/g}$, wet weight)

Lake and Species	Mean	Min	Max
<u>Baptiste Lake</u>			
Lake whitefish	0.19	0.12	0.23
Northern pike	0.35	0.23	0.64
Walleye	0.52	0.25	0.96
<u>Eagle Lake</u>			
Northern pike	0.07	0.05	0.09
Walleye	0.21	0.07	0.51
<u>Elnor Lake</u>			
Lake whitefish	0.11	0.02	0.27
Northern pike	0.70	0.32	1.89
Walleye	0.73	0.23	1.59
<u>Ethel Lake</u>			
Lake whitefish	0.04	0.02	0.08
Northern pike	0.42	0.10	0.61
Walleye	0.45	0.24	0.89
<u>Goodfish Lake</u>			
Northern pike	0.46	0.27	0.58
Walleye	0.28	0.17	0.45
<u>Greene Lake</u>			
Lake whitefish	0.05	0.03	0.06
Walleye	0.19	0.13	0.27
<u>Hilda Lake</u>			
Northern pike	0.50	0.31	0.79
Walleye	0.67	0.34	0.98
<u>Kaho Lake</u>			
Lake whitefish	0.11	0.04	0.39
Northern pike	0.14	0.09	0.26
Walleye	0.22	0.14	0.59
<u>Lac la Poudre</u>			
Lake whitefish	0.07	0.03	0.23
Northern pike	0.31	0.19	0.43
Walleye	0.55	0.20	1.17
Yellow perch	0.08	0.07	0.09
<u>Lac Ste Anne</u>			
Walleye	0.14	0.04	0.24
<u>Lake Isle</u>			
Northern pike	0.07	0.06	0.09
<u>Marie Lake</u>			
Lake whitefish	0.04	0.02	0.16
Northern pike	0.24	0.08	0.41
Walleye	0.19	0.10	0.37
<u>McLeod Lake</u>			
Rainbow Trout	0.05	0.03	0.07
<u>Moonshine Lake</u>			
Rainbow Trout	0.07	0.04	0.12
<u>Moose Lake</u>			
Cisco	0.09	0.04	0.16
Lake whitefish	0.06	0.03	0.11
Northern pike	0.38	0.11	1.32
Walleye	0.49	0.18	0.77
Yellow perch	0.05	0.00	0.12

Lake and Species	Mean	Min	Max
<u>Pigeon Lake</u>			
Lake whitefish	0.04	0.02	0.06
Walleye	0.14	0.05	0.28
<u>Pine Lake</u>			
Northern pike	0.22	0.14	0.34
Walleye	0.27	0.12	0.66
<u>Rock Island Lake</u>			
Northern pike	0.08	0.03	0.33
Walleye	0.11	0.04	0.30
<u>Sylvan Lake</u>			
Lake whitefish	0.12	0.03	0.22
Walleye	0.21	0.10	0.64
<u>Whitefish Lake</u>			
Lake whitefish	0.12	0.01	0.32
Northern pike	0.66	0.46	0.83
Walleye	0.78	0.41	1.30
<u>Wizard Lake</u>			
Northern pike	0.20	0.10	0.35




Figure 6 Mean Concentrations of Total Mercury in Fish, 2012/2013

Mean THg concentrations in walleye, northern pike and lake whitefish collected in previous studies from other water bodies of Alberta are listed in Table 10 (AHW 2009a, b, c, d). There have been recent concerns raised about mercury releases from oil sands development leading to increasing mercury levels in fish in the Athabasca River ecosystem. Evans and Talbot (2012) performed an extensive analysis of data for mercury in walleye, northern pike and lake whitefish. For the Athabasca River (1984-2003) they found decreasing trends for mercury in walleye and lake whitefish. For western Lake Athabasca and its delta, they found a decreasing trend for northern pike (1981-2009) and no trend for walleye (1981-2005). A recent study on mercury deposition in the Athabasca region has shown that there is an increase in mercury concentrations near oil sands operations (Kirk et al. 2014). The increased concentration could impact aquatic ecosystems.

Mean THg concentrations determined by Environment Canada for Lake Athabasca from 2010 to 2013 were consistent with the results obtained by Departments of Environment and Sustainable Resource Development and Health (Appendix A).

Mean THg concentrations in walleye collected in Rolling Hills Reservoir and Touchwood Lake in 2009/2010, in Amisk Lake, Kinnaird Lake, and Pinehurst Lake in 2010/2011, and in Elinor Lake, Hilda Lake, and Whitefish Lake in 2012/2013 were higher than those collected in other water bodies of Alberta previously.

Mean THg concentrations in northern pike collected in McGregor Lake, Rolling Hills Reservoir, Touchwood Lake and Whitefish Lake in 2009/2010, in Amisk Lake and Pinehurst Lake in 2010/2011 and in Elinor Lake and Whitefish Lake in 2012/2013 were higher than those collected in other water bodies of Alberta previously.

Mean THg concentrations in lake whitefish collected in Whitefish Lake in 2009/2010 were higher than those collected in other water bodies of Alberta previously.

The average THg concentrations in Canadian market fish reported by Health Canada ranged from 0.02 to 1.82 µg/g (Health Canada 2007). Compared to Canadian market fish for different fish species, mean THg concentrations in local fish were within the range of Canadian market fish.

Mean THg levels for walleye and northern pike from other water bodies in Canada and the U.S. reported in the literature are summarized in Table 11. Mean THg concentrations for the same fish species collected in the water bodies of Alberta in 2009/2013 were well within the ranges for the same fish species reported in the literature for other North American freshwater fish.

Mean THg concentration in fish fillets varied in other lakes, rivers and reservoirs in Canada and the U.S. The highest mean mercury levels in walleye and Northern pike in the water bodies in eastern and northern Canada ranged from 3.73 to 6.44 µg/g, respectively. High levels tended to be found in larger, older fish. Fish absorb MeHg directly through their gills or through the consumption of prey which contain mercury.

MeHg is tightly bound to proteins in all fish tissue so larger, older fish contain higher mercury (Munn and Short 1997, Neumann and Ward 1999).

Trophic level is a major factor in mercury accumulation in predatory (fish-eating) fish through biomagnification (Cabana et al. 1994). Bottom-feeding species may accumulate high mercury concentrations from direct contact with contaminated sediment or by eating benthic invertebrates and epibenthic organisms. Predatory fish species may accumulate and biomagnify mercury concentrations via several trophic levels of the food web (Suedel et al. 1994). Predators are commonly used as good indicators of mercury contamination. In this survey, the higher mercury levels were observed in walleye and northern pike which are highly piscivorous predatory fish.

Table 10 Mean THg Concentrations in Fish Muscles in Alberta Water Bodies

	Mean (µg/g, ww)	Location	Reference
Walleye	0.28 – 0.47	Athabasca River	AHW 2009a, 2009b, 2009c, 2009d
	0.42	Christina Lake	
	0.30	Clearwater River	
	0.13 – 0.16	Gregoire Lake	
	0.27	Keho Lake	
	0.43	Milk River Ridge Reservoir	
	0.63	Lac la Nonne	
	0.13	Lac Ste. Anne	
	0.41	Lake Newell	
	0.11	Pigeon Lake	
	0.52 – 0.79	Pine Coulee Reservoir	
	0.41	Pine Lake	
	0.79	Red Deer River	
	0.68	South Saskatchewan River	
	0.13	Winefred Lake	
Northern pike	0.42	Christina Lake	AHW 2009a, 2009b, 2009c, 2009d
	0.15 – 0.30	Clearwater River	
	0.15 - 0.21	Gregoire Lake	
	0.22	Keho Lake	
	0.56	Lac la Nonne	
	0.14	Lac Ste. Anne	
	0.23	Lake Newell	
	0.27 – 0.59	Little Bow River downstream	
	0.04 – 0.29	Little Bow River downstream	
	0.21	Milk River Ridge Reservoir	
	0.11 - 0.22	Muskeg River	
	0.18	Oldman River	
	0.13	Pine Coulee Reservoir	
	0.27	Pine lake	
	0.27	Red Deer River	
	0.35	South Saskatchewan River	
	0.44 – 0.56	Twin Valley Reservoir	
	0.27 – 0.49	Willow Creek Downstream	
	0.08 – 0.13	Winefred Lake	
Lake whitefish	0.09 – 0.17	Athabasca River	AHW 2009a, 2009b, 2009c, 2009d
	0.09	Christina Lake	
	0.04	Gregoire Lake	
	0.10	Keho Lake	
	0.13	Lake Newell	
	0.14	Milk River Ridge Reservoir	
	0.02	Pigeon Lake	
	0.13	South Saskatchewan River	
	0.08	Winefred Lake	

Table 11 Mean THg Levels in Fish Muscles Reported in the Literature

Species	Mean (µg/g, ww)	Location	Reference
Walleye	0.5	Northwest Ontario lakes (used as a reference for Walleye system)	Neff et al. 2012
	0.15 – 0.45	Lake Ontario (seasonal differences)	Zhang et al. 2012
	0.078 – 2.3 dry weight	Boreal lakes within 107 km of Sudbury	Yang et al. 2010
	0.15 to 0.6	Great Lakes with Lake Erie lowest to Lake Superior highest	Bhavsar et al. 2010
	0.05 – 0.90	16 Lakes, Northern Glaciated Plains, US	Setch et al. 2007
	0.19 – 0.30	Reservoirs, Manitoba, Canada	Bixley et al. 2007
	0.42 – 2.96	Walleye River system, Ontario	Kinghorn et al. 2007
	0.89 – 1.00	19 undisturbed lakes, Haute Mauricie, Quebec, Canada	Garcia and Carignan, 2005
	1.29 – 3.73	18 disturbed lakes, Haute Mauricie, Quebec, Canada	Garcia and Carignan, 2005
	0.750	Water bodies in northeastern of US and Canada (N=19,178)	Kamman et al. 2005
	0.58	Great Lakes, US	Gerstenberger and Dellinger, 2002
	0.47	Lakes in Northern Canada	Lockhart et al. 2005
	0.05 – 1.34	Canadian Arctic, Canada	Braune et al. 1999
	0.32 – 1.26	29 Lakes in the La Grande complex watershed, Quebec, Canada	Vendon et al. 1991
	0.19 – 1.43	Mackenzie River Basin Lakes	Evans et al. 2005a
Northern pike	0.4	Northwest Ontario lakes (used as a reference for Walleye system)	Neff et al. 2012
	0.2 – 0.35	Lake Ontario (seasonal differences)	Zhang et al. 2012
	0.5	Twin Valley Reservoir, southern Alberta: 2 year old	Brinkmann & Rasmussen 2010
	1.22	Twin Valley Reservoir, southern Alberta: 5 - 6 year old	Brinkmann & Rasmussen 2010
	0.2 – 0.23	Great Lakes, Lake Huron lowest to Lake Superior highest	Bhavsar et al. 2010
	0.26 – 0.32	Reservoirs, Manitoba, Canada	Bixley et al. 2007
	0.44 – 2.14	Walleye River system, Ontario, Canada	Kinghorn et al. 2007
	1.00 – 2.55	19 undisturbed lakes, Haute Mauricie, Quebec, Canada	Garcia and Carignan, 2005
	1.90 – 6.44	18 disturbed lakes, Haute Mauricie, Quebec, Canada	Garcia and Carignan, 2005
	0.045	lakes, rivers and reservoirs in northeastern of US and Canada (N=19,178)	Kamman et al. 2005
	0.16 – 1.1	Mackenzie River Basin, Canada	Evans, et al. 2005a
	0.12 – 0.74	Mackenzie River Basin, Canada	Evans, et al. 2005b
	0.378	Lakes in Northern Canada	Lockhart et al. 2005
	0.623 – 1.51	Yukon River, Kuskokwim River, US	Jewett et al. 2003
	0.11 – 0.63	Canadian Arctic, Canada	Braune et al. 1999
	0.25 – 0.90	29 Lakes in the La Grande complex watershed, Quebec, Canada	Vendon et al. 1991

Lake elevation	< 0.1	Northwest Ontario lakes listed as a reference for Atabasco system	Mad et al. 2012
	0.05 – 0.07	Reservoirs, Manitoba, Canada	Stetler et al. 2007
	0.08 – 0.11	Atabasco River system, Ontario	Kinghorn et al. 2007
	0.14 – 1.16	12 undisturbed lakes, Haute Mauricie, Quebec, Canada	Garnier and Carignan, 2005
	0.11 – 1.16	12 disturbed lakes, Haute Mauricie, Quebec, Canada	Garnier and Carignan, 2005
	0.209	Water bodies at northwestern of US and Canada (N=19, 17%)	Karickhoff et al. 2005
	0.01	Great Lakes, US	Gendronberger and Dettling, 2002
	0.14 – 0.16	MacKenzie River Basin, Canada	Fox et al. 2005
	0.11 – 0.13	Lakes in Northern Canada	Lockhart et al. 2005
	0.02 – 0.02	Canadian Arctic, Canada	Brauer et al. 1999
	0.07 – 0.16	26 Lakes in the La Grande complex watershed, Quebec, Canada	Verrier et al. 1991

3.2 Local Fish Consumption Rates

Three surveys of fish consumption patterns were conducted in communities of Central Alberta between 1997 and 2000. The first survey was conducted by Alberta Health and Wellness in Swan Hills communities in 1997 (AHW 1997). The second survey was conducted by the First Nations and Inuit Health Branch (FNIH) of Health Canada for the First Nations people living in the Lesser Slave Lake area in 1999 (Health Canada 1999). The third survey was conducted by the Environmental Health Sciences Program at the University of Alberta for the residents living in the communities near the Athabasca River and tributaries at Hinton (EHSUA 2000).

Table 12 Local Fish Consumption Rates in Communities of Central Alberta

Group	Subsistence Consumer Lesser Slave Lake ^a mean (g/d) % ^{**} (n=125)		Local Fish Consumer Swan Hills mean (g/d) % (n=127)		Local Fish Consumer Athabasca River mean (g/d) % (n=45)	
High (>100g/d)	273	5	107	2	121	2
Medium (30-99 g/d)	45	14	47	13	51	6
Low (5-29 g/d)	13	28	13	28	15	28
Very Low (≤ 4g/d)	1.6	43	2	57	1.6	66

^a mean from Phase I and Phase II studies (Health Canada 1999). ^{**} % of surveyed population

Fish consumption rates in different ethnic groups from these surveys are summarized in Table 12. A small proportion of local anglers and First Nations people consumed total fish over 100 g/d. Five per cent of the First Nations people in the Lesser Slave Lake communities were high consumers who ate total fish at an average of 273 g/d, much higher than the 2% of those in Beaver Hills communities who were high consumers at an average of 167 g/d and those in the communities nearby Hinton who were high consumers at an average rate of 129 g/d. The total fish consumption rates in the survey of the Lesser Slave Lake were similar with the results of the Beaver Hills survey in medium, low and very low intake groups. The majority of total fish consumers (85%, 80%) consumed fish at a low rate of 1.3 – 15 g/d. The majority of the First Nations group (87%) consumed fish at a low rate of 1.3 – 13 g/d.

The most common fish species consumed by the surveyed populations were northern pike, walleye, lake whitefish, and lake trout by the First Nations people in the Lesser Slave Lake communities, walleye, northern pike, perch, brook trout, Lake whitefish and Arctic grayling by the residents in Beaver Hills communities, and northern pike, Arctic grayling, mountain whitefish, northern pike and walleye by the residents in the communities nearby Hinton.

The results from the above surveys were derived from adults only. Fish consumption rates (g/d) vary in different subpopulations (IJC/PA 2000). Children may consume larger quantities compared to their body weight than adults. Prenatal exposure may occur through placental transfer. For the purpose of risk management, these subpopulations are considered as potential high risk groups for exposure to mercury from fish consumption.

3.3 Estimated Exposures

Exposure ratios were estimated for consuming walleye and northern pike. Estimated exposure ratios based on the estimated fish consumption derived by the gTUEs from Inuit in Canada are summarized in Table 13 for women of reproductive age and Table 14 for adults. Specific fish consumption rates were not available for women of reproductive age and young children. As a result, the estimation of exposures for young children was not performed. The fish consumption rate for all adults was used for estimating exposures for women of reproductive age.

In general, the estimated exposure ratios were greater than one for the high intake group, especially for a subpopulation of women of reproductive age if consuming predatory fish like the large walleye and northern pike. The values of gTUEs were derived from risk assessment approaches with many assumptions and uncertainties. The risk assessment is specifically designed to avoid underestimating risk. The results do not mean that specific individuals or populations have immediate or even their health consequences from mercury exposure. An estimated exposure ratio greater than one should be used as a reference point for making risk management decisions. In particular, those exposure scenarios with an exposure ratio greater than one warrant closer attention including providing information about maximum recommended fish consumption to allow consumers to make safe eating choices.

Many factors influence the estimated population density such as body weight and consumption rates. The body weight of 15 kg used in this assessment was derived from the 1988 National Population Health Survey in Alaska adults. In this report, the age-specific body weights for women of reproductive age and young children in Alaska were not available. The average body weights (adults) (adults) were 15 kg for women of reproductive age, 15 kg for the male 15-24 age group, and 15 kg for the male 25-34 age group. The consumption rates used in this report were based on the three surveys of adults living in Central Alaska. The estimated population was solely based on data from these specific surveys. People may also be interested in knowing their impact on the environment and other natural resources.

[illegible][illegible]

Year	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100
1990	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100

Lake and Species	Lead Contaminant High Intake (175 µg/d)	Lead Contaminant Medium Intake (50 µg/d)	Substance Contaminant High Intake (275 µg/d)
<u>North Malensee Lake</u>			
Northern pike	6.1	1.2	6.6
Mudminnow	3.5	1.0	3.6
<u>Portage Lake</u>			
Lake trout	1.9	< 1	3.0
<u>Plover Lake</u>			
Lake whitefish	< 1	< 1	< 1
Mudminnow	1.9	< 1	3.0
<u>Pine Lake</u>			
Northern pike	2.9	< 1	6.6
Mudminnow	3.6	1.1	5.7
<u>Portland Lake</u>			
Northern pike	11	3.1	17
Mudminnow	7.9	2.3	13
<u>Reuben Lake</u>			
Lake trout	9.0	2.9	16
<u>Rutherford Lake</u>			
Northern pike	3.7	1.1	5.9
Mudminnow	3.0	< 1	4.8
Lake whitefish	1.0	< 1	1.5
<u>Rock Island Lake</u>			
Northern Pike	1.0	< 1	1.6
Mudminnow	1.0	< 1	2.3
<u>Riding Lake Reservoir</u>			
Northern pike	16	4.0	22
Mudminnow	10	4.3	26
<u>Shanley Lake</u>			
Northern pike	1.6	1.0	2.5
Mudminnow	2.1	1.0	3.3
<u>Snake Lake</u>			
Northern pike	0.7	< 1	1.0
Mudminnow	0.5	< 1	1.0
<u>Sturgeon Lake</u>			
Lake whitefish	< 1	< 1	< 1
Northern pike	1.0	< 1	3.1
Mudminnow	2.3	< 1	3.8
<u>Swan Lake</u>			
Lake whitefish	1.5	< 1	2.6
Mudminnow	2.7	< 1	4.6
<u>Tweed Lake</u>			
Lake whitefish	1.6	< 1	2.3
Northern pike	11	3.3	18.9
Mudminnow	12	3.6	20
<u>Wabigoon Lake</u>			
Lake whitefish	0.40	0.13	0.73
Northern pike	6.30	1.67	7.91
<u>Whitfish Lake (2009/2010)</u>			
Lake whitefish	9.6	2.8	15.0
Northern pike	6.6	2.5	13
Mudminnow	11	3.3	18

Lake and Species	Local Consumer High Intake (170 g/d)	Local Consumer Medium Intake (50 g/d)	Subsistence Consumer High Intake (270 g/d)
<u>Whitefish Lake (2012/2013)^a</u>			
Lake whitefish	1.6	< 1	2.6
Northern pike	8.6	2.5	13.7
Walleye	10.1	3.0	16.1
<u>Winnipeg Lake</u>			
Northern pike	1.6	1.0	2.5
Walleye	2.5	1.0	4.0
<u>Wizard Lake</u>			
Northern pike	2.6	< 1	4.1
<u>Wolf Lake</u>			
Lake whitefish	1.4	< 1	2.1
Northern pike	4.4	1.3	7.0
Walleye	7.7	2.3	12.3

Note: mean of total mercury listed in Tables 6, 7, 8, and 9, body weight = 65 kg, pTDI = 0.2 µg/kg bw/d for women

^a Exposure ratios are rounded to 2 significant figures in keeping with the precision of the data they are based upon. ^b Sampling year for lake if lake was sampled in different years. ^c calculations based on data from Environment Canada

Table 14 Estimated Exposure Ratios for Adults

Lake and Species	Local Consumer High Intake (170 g/d)	Local Consumer Medium Intake (50 g/d)	Subsistence Consumer High Intake (270 g/d)
<u>Amisk Lake</u>			
Northern pike	2.7	1.0	4.3
Walleye	4.4	1.3	7.0
<u>Baptiste Lake (2011/2012)^a</u>			
Northern pike	< 1	< 1	1.1
Walleye	< 1	< 1	< 1
<u>Baptiste Lake (2012/2013)^a</u>			
Lake	< 1	< 1	1.5
whitefish			
Northern pike	1.7	< 1	2.7
Walleye	2.6	< 1	4.1
<u>Beaver Lake</u>			
Lake whitefish	< 1	< 1	< 1
Northern pike	2.2	< 1	3.6
Walleye	1.8	< 1	2.8
<u>Bitscho Lake</u>			
Northern pike	< 1	< 1	0.6
Walleye	1.0	< 1	0.9
<u>Bourque Lake</u>			
Lake whitefish	< 1	< 1	< 1
Northern pike	1.5	< 1	2.3
Walleye	1.8	< 1	2.8
<u>Calina Lake</u>			
Northern pike	< 1	< 1	1.3
Walleye	< 1	< 1	1.2
<u>Chinchaqua River</u>			
Walleye	2.2	1.0	3.5
<u>Chrystina Lake</u>			
Brook trout	1.0	< 1	1.6
<u>Cowoki Reservoir</u>			
Northern pike	2.9	< 1	4.6
Walleye	2.3	< 1	3.7
<u>Crawling Valley Reservoir</u>			
Northern pike	1.6	< 1	3.0
Walleye	2.6	< 1	4.2
<u>Cross (Steele) Lake</u>			
Northern pike	1.0	< 1	1.7
<u>Dore Lake^b</u>			
Northern pike	< 1	< 1	1.3
Walleye	1.1	< 1	1.7
<u>Enoch Lake</u>			
Northern pike	< 1	< 1	< 1
Walleye	1.0	< 1	1.6
<u>Edith Lake</u>			
Brook trout	< 1	< 1	1.4
<u>Elinor Lake</u>			
Lake whitefish	< 1	< 1	< 1
Northern pike	3.5	1.0	5.5
Walleye	3.6	1.1	5.7

Lake and Species	Local Consumer High Intake (170 g/d)	Local Consumer Medium Intake (50 g/d)	Subsistence Consumer High Intake (270 g/d)
<u>Ethel Lake</u>			
Lake whitefish	< 1	< 1	< 1
Northern pike	2.1	< 1	3.3
Walleye	2.2	< 1	3.6
<u>Goodfish Lake</u>			
Northern pike	2.3	< 1	3.6
Walleye	1.4	< 1	2.2
<u>Gregoire Lake</u>			
Lake whitefish	< 1	< 1	< 1
Walleye	< 1	< 1	1.5
<u>Hay River</u>			
Walleye	3.0	< 1	4.8
<u>Heart Lake</u>			
Lake whitefish	< 1	< 1	< 1
Northern pike	< 1	< 1	1.2
Walleye	< 1	< 1	< 1
<u>Hilda Lake</u>			
Northern pike	2.5	< 1	3.9
Walleye	3.3	< 1	5.3
<u>Hutch Lake</u>			
Walleye	< 1	< 1	< 1
<u>Isle Lake (2009/2010)^a</u>			
Northern pike	< 1	< 1	< 1
Walleye	1.0	< 1	1.6
<u>Isle Lake (2010/2011)^a</u>			
Northern pike	1.9	1.0	3.0
Lake whitefish	< 1	< 1	< 1
<u>Kehewin Lake</u>			
Northern pike	1.9	< 1	3.1
Walleye	1.4	< 1	2.3
<u>Keho Lake</u>			
Lake whitefish	< 1	< 1	< 1
Northern pike	< 1	< 1	1.1
Walleye	1.1	< 1	1.7
<u>Kinnaird Lake</u>			
Northern pike	2.1	1.0	3.4
Walleye	3.3	1.0	5.3
<u>Kirby Lake</u>			
Lake whitefish	< 1	< 1	< 1
Northern pike	< 1	< 1	< 1
<u>Lac Bellevue</u>			
Walleye	1.7	1.0	2.8
<u>Lac la Nonne</u>			
Lake whitefish	< 1	< 1	< 1
Northern pike	1.5	< 1	2.5
Walleye	2.7	< 1	4.3
Yellow Perch	< 1	< 1	< 1
<u>Lac Ste Anne</u>			
Walleye	< 1	< 1	1.1

Lake and Species	Local Consumer High Intake (170 g/d)	Local Consumer Medium Intake (50 g/d)	Subsistence Consumer High Intake (270 g/d)
<u>Lake Athabasca</u>			
Lake trout	1.3	< 1	2.1
Lake whitefish	< 1	< 1	< 1
Northern pike	1.0	< 1	1.7
Walleye	1.3	< 1	2.0
<u>Lake Isle (2012/2013)^a</u>			
Northern Pike	< 1	< 1	< 1
<u>Laurier Lake</u>			
Northern pike	2.3	1.0	3.7
<u>Len Thompson</u>			
Rainbow Trout	1.0	< 1	1.1
<u>Lesser Slave Lake East Basin</u>			
Northern pike	2.0	1.0	3.2
Walleye	1.6	1.0	2.6
<u>Lesser Slave Lake West Basin</u>			
Northern pike	1.6	1.0	2.6
Walleye	1.7	1.0	2.8
<u>Little Bow Reservoir</u>			
Northern pike	2.3	< 1	3.7
<u>Long Lake</u>			
Northern pike	1.7	1.0	2.7
Walleye	2.1	1.0	3.4
<u>Loon River</u>			
Walleye	1.7	< 1	2.6
<u>Marie Lake</u>			
Lake whitefish	< 1	< 1	< 1
Northern pike	1.2	< 1	1.9
Walleye	< 1	< 1	1.5
<u>McGregor Lake</u>			
Northern pike	5.0	1.5	7.9
Walleye	3.1	< 1	4.9
<u>McLeod Lake</u>			
Rainbow Trout	< 1	< 1	< 1
<u>McMillan Lake</u>			
Northern pike	1.9	< 1	3.1
<u>Meander River</u>			
Northern pike	1.0	< 1	1.5
Walleye	1.0	< 1	1.3
<u>Moonshine Lake</u>			
Rainbow Trout	< 1	< 1	< 1
<u>Moose Lake (2011/2012)^a</u>			
Northern pike	1.6	< 1	2.6
Walleye	2.4	< 1	3.9
<u>Moose Lake (2012/2013)^a</u>			
Cisco	< 1	< 1	< 1
Lake whitefish	< 1	< 1	< 1
Northern pike	< 1	< 1	< 1
Walleye	< 1	< 1	1.4
Yellow Perch	< 1	< 1	< 1

Lake and Species	Local Consumer High Intake (170 g/d)	Local Consumer Medium Intake (50 g/d)	Subsistence Consumer High Intake (270 g/d)
<u>Nipisi Lake</u>			
Northern pike	< 1	< 1	< 1
<u>North Wabasca Lake</u>			
Lake Whitefish	3.6	1.0	5.7
Northern pike	1.5	1.0	2.4
Walleye	1.3	< 1	2.1
<u>Peerless Lake</u>			
Lake trout	< 1	< 1	1.1
<u>Pigeon Lake</u>			
Lake whitefish	< 1	< 1	< 1
Walleye	< 1	< 1	1.1
<u>Pine Lake</u>			
Northern pike	1.1	< 1	1.8
Walleye	1.4	< 1	2.2
<u>Pinehurst Lake</u>			
Northern pike	4.0	1.2	6.4
Walleye	3.0	1.0	4.7
<u>Pitchimi Lake</u>			
Lake trout	3.7	1.1	5.9
<u>Richardson Lake</u>			
Northern pike ^b	1.4	< 1	2.2
Walleye ^b	1.1	< 1	1.8
Lake whitefish	< 1	< 1	1.0
<u>Rock Island Lake</u>			
Northern pike	1.0	< 1	1.6
Walleye	1.5	< 1	2.3
<u>Rolling Hills Reservoir</u>			
Northern pike	5.2	1.5	8.2
Walleye	5.6	1.6	8.9
<u>Skeleton Lake</u>			
Northern pike	1.0	< 1	1.0
Walleye	1.0	< 1	1.3
<u>Snipe Lake</u>			
Northern pike	< 1	< 1	< 1
Walleye	< 1	< 1	< 1
<u>Sturgeon Lake</u>			
Lake whitefish	< 1	< 1	< 1
Northern pike	< 1	< 1	1.2
Walleye	< 1	< 1	1.4
<u>Sylvan Lake</u>			
Lake whitefish	< 1	< 1	< 1
Walleye	1.0	< 1	1.7
<u>Touchwood Lake</u>			
Lake whitefish	< 1	< 1	< 1
Northern pike	4.3	1.2	6.8
Walleye	4.7	1.3	7.4
<u>Wabamun Lake</u>			
Lake whitefish	0.17	< 1	0.28
Northern pike	1.89	1.0	3.0

Lake and Species	Local Consumer High Intake (170 g/d)	Local Consumer Medium Intake (50 g/d)	Subsistence Consumer High Intake (270 g/d)
<u>Whitefish Lake (2009/2010)^a</u>			
Lake whitefish	3.6	1.0	5.7
Northern pike	3.2	< 1	5.0
Walleye	4.3	1.3	6.8
<u>Whitefish Lake (2012/2013)^a</u>			
Lake whitefish	< 1	< 1	< 1
Northern pike	3.3	< 1	5.2
Walleye	3.8	1.1	6.1
<u>Winigami Lake</u>			
Northern pike	1.0	< 1	1.0
Walleye	1.0	< 1	1.5
<u>Wizard Lake</u>			
Northern pike	< 1	< 1	1.5
<u>Wolf Lake</u>			
Lake whitefish	< 1	< 1	< 1
Northern pike	1.7	< 1	2.7
Walleye	2.9	< 1	4.7

Note: mean of total mercury listed in Tables 6, 7, 8 and 9; body weight = 73 kg; pTDI = 0.47 µg/kg bw/d for adults.

^a Sampling year for lake if lake was sampled in different years. ^b calculations based on data from Environment Canada

3.4 Consumption Limits

For the purpose of quantitative fish advisories, the lifetime consumption limits were calculated for subgroups of women, young children and adults if total mercury levels in fish exceeded 0.2 µg/g (Table 15).

These consumption limits were specific to fish species and site. The values provide the information on the maximum amount of local fish that can be safely consumed on a weekly basis for a lifetime by the identified subpopulations for fish specified from the specified locations. Fish preparation and cooking methods do not reduce the concentrations of total mercury in fish (Morgan et al. 1997).

Table 15 Lifetime Fish Consumption Limits^a

Lake and Species	Thg µg/g	Women g/week	Children (5-11 yr) g/week	Children (1-4 yr) g/week	Adults g/week
Arrow Lake					
Northern pike	0.54	170	68	37	660
Walleye	0.89	100	42	23	270
Badlake Lake					
Northern pike	0.38	240	97	53	633
Walleye	0.52	176	72	39	465
Beaver Lake					
Northern pike	0.45	201	82	44	530
Walleye	0.35	258	105	57	680
Bowman Lake					
Northern pike	0.30	305	124	67	804
Walleye	0.38	253	103	56	667
Chinook River					
Walleye	0.44	210	84	46	550
Conch Reservoir					
Northern pike	0.59	150	63	34	410
Walleye	0.47	180	79	41	490
Crowfoot Valley Reservoir					
Northern pike	0.38	240	97	53	630
Walleye	0.54	170	68	37	440
Cross (Steep) Lake					
Northern pike	0.21	430	174	95	1134
Dore Lake^c					
Northern pike	0.34	260	109	59	707
Walleye	7/57	150	65	35	420
Ends Lake					
Walleye	0.21	430	178	97	1157
Elmer Lake					
Northern pike	0.70	130	53	29	343
Walleye	0.73	125	51	28	329
Elton Lake					
Northern pike	0.42	218	89	48	579
Walleye	0.45	201	82	44	531
Goodfish Lake					
Northern pike	0.46	198	81	44	524
Walleye	0.28	324	131	72	854
Hay River					
Walleye	0.61	150	61	33	400
Hole Lake					
Northern pike	0.50	182	74	40	481
Walleye	0.67	138	55	30	358
Isle Lake					
Northern pike	0.38	240	97	53	630
Refinery Lake					
Northern pike	0.39	230	95	52	620
Walleye	0.29	310	120	70	830
Rain Lake					
Walleye	0.22	410	168	91	1081

Lake and Species	Tkg 10/5	Sturgeon g/mesh	Children (5-11 yr) g/mesh	Children (1-4 yr) g/mesh	Adults g/mesh
Canard Lake					
Northern pike	0.44	210	50	40	100
Sturgeon	0.47	140	30	30	80
Laç Joliveau					
Sturgeon	0.35	200	100	20	400
Laç à Marie					
Northern pike	0.31	200	110	60	710
Sturgeon	0.35	100	67	37	437
Lake Abitibi					
Lake trout	0.27	100	100	70	800
Northern pike	0.31	410	180	40	1,100
Sturgeon	0.25	300	140	70	820
Lac de la					
Sturgeon	0.47	100	70	40	310
Lower Saint-Louis East Basin					
Northern pike	0.33	200	110	61	730
Sturgeon	0.35	200	110	50	600
Lower Saint-Louis West Basin					
Northern pike	0.40	230	80	50	630
Sturgeon	0.33	200	110	61	730
Lake Bow Reservoir					
Northern pike	0.47	100	70	40	310
Lake Lake					
Northern pike	0.30	270	110	40	710
Sturgeon	0.43	270	60	67	380
Lake River					
Sturgeon	0.34	271	110	60	710
Le Lac					
Northern pike	0.24	370	180	90	900
McGarry Lake					
Northern pike	1.00	81	37	20	200
Sturgeon	0.40	100	60	30	300
McMillan Lake					
Northern pike	0.30	230	20	60	610
Moore Lake					
Northern pike	0.30	340	60	50	600
Sturgeon	0.40	100	70	61	600
North Muskegon Lake					
Northern pike	0.31	240	130	60	710
Sturgeon	0.27	340	140	70	800
Pine Lake					
Northern pike	0.33	400	180	50	1070
Sturgeon	0.27	330	130	70	810
Pinehurst Lake					
Northern pike	0.51	110	40	20	300
Sturgeon	0.60	100	60	30	400
Robins Lake					
Lake trout	0.75	120	60	27	320
Richardson Lake					
Northern pike	0.26	320	130	71	840
Sturgeon	0.25	380	180	87	1020
Sturgeon	1.1	81	30	18	210

Lake and Species	Age range	Mean growth	Outlier 0.47 at growth	Outlier 0.47 at growth	Adult growth
Yellow Lake, Walleye					
Northern age	1-5	80	80	10	100
Multiple	1-1	81	85	8	100
Yellow Lake, Walleye					
Northern age	0-25	400	110	0	1100
Yellow Lake, Walleye					
Northern age	0-20	100	90	20	100
Multiple	0-20	87	90	21	100
Yellow Lake, Walleye					
Northern age	0-20	200	97	50	400
Yellow Lake, Walleye					
Northern age	0-10	100	90	10	100
Multiple	0-20	100	90	21	100
Yellow Lake, Walleye					
Northern age	0-20	80	100	100	100
Yellow Lake, Walleye					
Northern age	0-20	200	100	10	100
Multiple	0-20	100	90	10	100

Note: Mean growth = 1 kg for adults, 0.5 kg for juveniles. Outlier = 0.47 at growth. Adult = 1 kg for adults, 0.5 kg for juveniles. Multiple = 0.47 at growth for adults, 0.47 at growth for juveniles.

* Outlier = 0.47 at growth for adults, 0.47 at growth for juveniles. Outlier = 0.47 at growth for adults, 0.47 at growth for juveniles.

* Outlier = 0.47 at growth for adults, 0.47 at growth for juveniles.

3.3 Fish Consumption Advisories

Fish consumers may be exposed to mercury by consuming locally caught fish. Mercury is rapidly absorbed after ingestion and distributed throughout the body (WHO 1990). Mercury in the body is constantly excreted and can cross the placental and blood-brain barriers (Morgan et al. 1992). The half-life of mercury in the human body varies from 40 to 80 days (WHO 1990). Mercury leaves the human body via urine, feces and breast milk. Small amounts of ingested mercury are eliminated from the body with no marked adverse effects. At high exposure levels, mercury produces a variety of health effects. Larger amounts of mercury may damage the nervous system. Neurotoxicity may occur in the developing embryo or fetus during pregnancy and in young children. As a result, it is prudent to reduce mercury exposure for women of reproductive age and young children. The guidelines proposed by Health Canada are intended to protect susceptible populations.

Because mercury occurs naturally, mercury is found in all commercial fish. Commercial fish and other foods at low levels. People are exposed to very low levels of mercury via sources such as breathing the air, mercury amalgam dental fillings and eating other foods. Alberta Health and Wellness conducted a survey of mercury levels in blood, urine and hair in adults and children living in the Athabasca Lake and surrounding area communities in 2008 (AHWS 2008). The survey found that the

average levels of total mercury in blood, urine and hair in Chinese participants were lower than people living in other areas and countries.

Mercury levels are high enough to cause high exposure to certain types of fish and shellfish consumption of high consumption is associated. Although high consumption may not represent a consistent higher level of mercury, it may not have a consistent effect on mercury exposure. The results from these surveys from different studies indicate that total fish consumption is not the primary source of mercury exposure, rather it is more associated with mercury exposure.

In order to protect all human consumption, reducing a high consumption of mercury is not an appropriate option. High consumption of mercury is associated with mercury exposure. The results from these surveys from different studies indicate that total fish consumption is not the primary source of mercury exposure, rather it is more associated with mercury exposure. The results from these surveys from different studies indicate that total fish consumption is not the primary source of mercury exposure, rather it is more associated with mercury exposure.

Since the early 1980s, some fish consumption advisories related to mercury have been issued and published in the United States. In 1990, the government issued a advisory to the public, as the total fish consumption is not the primary source of mercury exposure, rather it is more associated with mercury exposure. The results from these surveys from different studies indicate that total fish consumption is not the primary source of mercury exposure, rather it is more associated with mercury exposure.

3.2 Benefits of Fish Consumption

The benefits and risk of fish consumption is a source of public health concern. Fish is an important source of nutrients for people. However, it contains harmful substances like the long-chain omega-3 fatty acids, the omega-6 fatty acids, and the omega-3 fatty acids. The results from these surveys from different studies indicate that total fish consumption is not the primary source of mercury exposure, rather it is more associated with mercury exposure.

Health Canada released the evidence showing an association between reduced risk of sudden cardiac death and fish consumption frequency. In 2007, Health Canada (2007) in a small case-control study, 79 cases, 100 controls, reported that a reduced risk of myocardial infarction with fish consumption of at least one meal per week was observed. The results from these surveys from different studies indicate that total fish consumption is not the primary source of mercury exposure, rather it is more associated with mercury exposure.

In contrast, a large number of people have been killed since 1987 and
in 1990. In addition, there has been a large number of people who have
been killed in the past few years. In the past few years, there have been
many people who have been killed in the past few years. In the past few
years, there have been many people who have been killed in the past few
years. In the past few years, there have been many people who have been
killed in the past few years. In the past few years, there have been many
people who have been killed in the past few years. In the past few years,
there have been many people who have been killed in the past few years.

The number of people who have been killed in the past few years is
very high. In the past few years, there have been many people who have
been killed in the past few years. In the past few years, there have been
many people who have been killed in the past few years. In the past few
years, there have been many people who have been killed in the past few
years. In the past few years, there have been many people who have been
killed in the past few years. In the past few years, there have been many
people who have been killed in the past few years. In the past few years,
there have been many people who have been killed in the past few years.

In the past few years, there have been many people who have been
killed in the past few years. In the past few years, there have been many
people who have been killed in the past few years. In the past few years,
there have been many people who have been killed in the past few years.

In the past few years, there have been many people who have been
killed in the past few years. In the past few years, there have been many
people who have been killed in the past few years. In the past few years,
there have been many people who have been killed in the past few years.

In the past few years, there have been many people who have been
killed in the past few years. In the past few years, there have been many
people who have been killed in the past few years. In the past few years,
there have been many people who have been killed in the past few years.

In the past few years, there have been many people who have been
killed in the past few years. In the past few years, there have been many
people who have been killed in the past few years. In the past few years,
there have been many people who have been killed in the past few years.

© 2005 Blackwell Publishing Ltd, *Journal of Internal Medicine* 258: 101–108

[illegible][illegible]

1987-1988-1989-1990-1991-1992-1993-1994-1995-1996-1997-1998-1999-2000-2001-2002-2003-2004-2005-2006-2007-2008-2009-2010-2011-2012-2013-2014-2015-2016-2017-2018-2019-2020-2021-2022-2023-2024-2025-2026-2027-2028-2029-2030-2031-2032-2033-2034-2035-2036-2037-2038-2039-2040-2041-2042-2043-2044-2045-2046-2047-2048-2049-2050-2051-2052-2053-2054-2055-2056-2057-2058-2059-2060-2061-2062-2063-2064-2065-2066-2067-2068-2069-2070-2071-2072-2073-2074-2075-2076-2077-2078-2079-2080-2081-2082-2083-2084-2085-2086-2087-2088-2089-2090-2091-2092-2093-2094-2095-2096-2097-2098-2099-2100-2101-2102-2103-2104-2105-2106-2107-2108-2109-2110-2111-2112-2113-2114-2115-2116-2117-2118-2119-2120-2121-2122-2123-2124-2125-2126-2127-2128-2129-2130-2131-2132-2133-2134-2135-2136-2137-2138-2139-2140-2141-2142-2143-2144-2145-2146-2147-2148-2149-2150-2151-2152-2153-2154-2155-2156-2157-2158-2159-2160-2161-2162-2163-2164-2165-2166-2167-2168-2169-2170-2171-2172-2173-2174-2175-2176-2177-2178-2179-2180-2181-2182-2183-2184-2185-2186-2187-2188-2189-2190-2191-2192-2193-2194-2195-2196-2197-2198-2199-2200-2201-2202-2203-2204-2205-2206-2207-2208-2209-2210-2211-2212-2213-2214-2215-2216-2217-2218-2219-2220-2221-2222-2223-2224-2225-2226-2227-2228-2229-2230-2231-2232-2233-2234-2235-2236-2237-2238-2239-2240-2241-2242-2243-2244-2245-2246-2247-2248-2249-2250-2251-2252-2253-2254-2255-2256-2257-2258-2259-2260-2261-2262-2263-2264-2265-2266-2267-2268-2269-2270-2271-2272-2273-2274-2275-2276-2277-2278-2279-2280-2281-2282-2283-2284-2285-2286-2287-2288-2289-2290-2291-2292-2293-2294-2295-2296-2297-2298-2299-2300-2301-2302-2303-2304-2305-2306-2307-2308-2309-2310-2311-2312-2313-2314-2315-2316-2317-2318-2319-2320-2321-2322-2323-2324-2325-2326-2327-2328-2329-2330-2331-2332-2333-2334-2335-2336-2337-2338-2339-2340-2341-2342-2343-2344-2345-2346-2347-2348-2349-2350-2351-2352-2353-2354-2355-2356-2357-2358-2359-2360-2361-2362-2363-2364-2365-2366-2367-2368-2369-2370-2371-2372-2373-2374-2375-2376-2377-2378-2379-2380-2381-2382-2383-2384-2385-2386-2387-2388-2389-2390-2391-2392-2393-2394-2395-2396-2397-2398-2399-2400-2401-2402-2403-2404-2405-2406-2407-2408-2409-2410-2411-2412-2413-2414-2415-2416-2417-2418-2419-2420-2421-2422-2423-2424-2425-2426-2427-2428-2429-2430-2431-2432-2433-2434-2435-2436-2437-2438-2439-2440-2441-2442-2443-2444-2445-2446-2447-2448-2449-2450-2451-2452-2453-2454-2455-2456-2457-2458-2459-2460-2461-2462-2463-2464-2465-2466-2467-2468-2469-2470-2471-2472-2473-2474-2475-2476-2477-2478-2479-2480-2481-2482-2483-2484-2485-2486-2487-2488-2489-2490-2491-2492-2493-2494-2495-2496-2497-2498-2499-2500-2501-2502-2503-2504-2505-2506-2507-2508-2509-2510-2511-2512-2513-2514-2515-2516-2517-2518-2519-2520-2521-2522-2523-2524-2525-2526-2527-2528-2529-2530-2531-2532-2533-2534-2535-2536-2537-2538-2539-2540-2541-2542-2543-2544-2545-2546-2547-2548-2549-2550-2551-2552-2553-2554-2555-2556-2557-2558-2559-2560-2561-2562-2563-2564-2565-2566-2567-2568-2569-2570-2571-2572-2573-2574-2575-2576-2577-2578-2579-2580-2581-2582-2583-2584-2585-2586-2587-2588-2589-2590-2591-2592-2593-2594-2595-2596-2597-2598-2599-2600-2601-2602-2603-2604-2605-2606-2607-2608-2609-2610-2611-2612-2613-2614-2615-2616-2617-2618-2619-2620-2621-2622-2623-2624-2625-2626-2627-2628-2629-2630-2631-2632-2633-2634-2635-2636-2637-2638-2639-2640-2641-2642-2643-2644-2645-2646-2647-2648-2649-2650-2651-2652-2653-2654-2655-2656-2657-2658-2659-2660-2661-2662-2663-2664-2665-2666-2667-2668-2669-2670-2671-2672-2673-2674-2675-2676-2677-2678-2679-2680-2681-2682-2683-2684-2685-2686-2687-2688-2689-2690-2691-2692-2693-2694-2695-2696-2697-2698-2699-2700-2701-2702-2703-2704-2705-2706-2707-2708-2709-2710-2711-2712-2713-2714-2715-2716-2717-2718-2719-2720-2721-2722-2723-2724-2725-2726-2727-2728-2729-2730-2731-2732-2733-2734-2735-2736-2737-2738-2739-2740-2741-2742-2743-2744-2745-2746-2747-2748-2749-2750-2751-2752-2753-2754-2755-2756-2757-2758-2759-2760-2761-2762-2763-2764-2765-2766-2767-2768-2769-2770-2771-2772-2773-2774-2775-2776-2777-2778-2779-2780-2781-2782-2783-2784-2785-2786-2787-2788-2789-2790-2791-2792-2793-2794-2795-2796-2797-2798-2799-2800-2801-2802-2803-2804-2805

doi:10.1017/S0022292412001606 Printed in the United Kingdom
© 2012 Cambridge University Press

1. *Erklären Sie, weshalb die folgenden Aussagen zutreffend sind:*
 a. Die Gewinnfunktion eines Unternehmens ist eine Funktion der eigenen und der fremden Produktionsmenge.
 b. Die Gewinnfunktion eines Unternehmens ist eine Funktion der eigenen und der fremden Produktionsmenge.
 c. Die Gewinnfunktion eines Unternehmens ist eine Funktion der eigenen und der fremden Produktionsmenge.
 d. Die Gewinnfunktion eines Unternehmens ist eine Funktion der eigenen und der fremden Produktionsmenge.

- a) Erklären Sie, was die Begriffe "Kultur" und "Werte" bedeuten und wie sie zusammenhängen.
- b) Nennen Sie drei Beispiele für kulturelle Unterschiede in der Wahrnehmung von Zeit und Raum.

6. Zusammenfassung

In der Vorlesung haben wir die Grundlagen der Kultur und der Werte kennen gelernt. Wir haben gesehen, dass Kultur ein breiter Begriff ist, der alle Aspekte des menschlichen Lebens umfasst. Werte sind die Prinzipien, nach denen wir handeln und die uns helfen, Entscheidungen zu treffen.

Wir haben auch gesehen, dass Kultur und Werte nicht statisch sind, sondern sich über die Zeit und über verschiedene Kulturen hinweg verändern. Dies ist ein Ergebnis der Interaktion zwischen verschiedenen Faktoren wie Technologie, Wirtschaft und Politik. Wir haben schließlich gesehen, dass Kultur und Werte eine wichtige Rolle in unserem Leben spielen und dass wir sie verstehen müssen, um die Welt um uns herum besser zu verstehen.

Wir haben gesehen, dass Kultur und Werte eine wichtige Rolle in unserem Leben spielen und dass wir sie verstehen müssen, um die Welt um uns herum besser zu verstehen. Wir haben gesehen, dass Kultur und Werte eine wichtige Rolle in unserem Leben spielen und dass wir sie verstehen müssen, um die Welt um uns herum besser zu verstehen.

Wir haben gesehen, dass Kultur und Werte eine wichtige Rolle in unserem Leben spielen und dass wir sie verstehen müssen, um die Welt um uns herum besser zu verstehen. Wir haben gesehen, dass Kultur und Werte eine wichtige Rolle in unserem Leben spielen und dass wir sie verstehen müssen, um die Welt um uns herum besser zu verstehen.

- a) Erklären Sie, was die Begriffe "Kultur" und "Werte" bedeuten und wie sie zusammenhängen.
- b) Nennen Sie drei Beispiele für kulturelle Unterschiede in der Wahrnehmung von Zeit und Raum.
- c) Diskutieren Sie, wie Kultur und Werte die Entwicklung einer Nation beeinflussen können.

Wir haben gesehen, dass Kultur und Werte eine wichtige Rolle in unserem Leben spielen und dass wir sie verstehen müssen, um die Welt um uns herum besser zu verstehen. Wir haben gesehen, dass Kultur und Werte eine wichtige Rolle in unserem Leben spielen und dass wir sie verstehen müssen, um die Welt um uns herum besser zu verstehen.

Table 15 Recommended Fish Consumption Limits

Water Body/Fish Species	Fish Size (lbs)	Consumption Limit (pounds per week)			
		Women	Children (6-11 yr)	Children (1-5 yr)	Adults
<u>Amisk Lake</u>					
Northern pike	2	avoid	avoid	avoid	0
Walleye	1	avoid	avoid	avoid	0
<u>Beartooth Lake</u>					
Northern pike	0	3	1	0.5	no limit
Walleye	2	avoid	avoid	avoid	0
<u>Bowyer Lake</u>					
Northern pike	5	3	1	0.5	no limit
Walleye	3	3	1	0.5	no limit
<u>Burns Lake</u>					
Northern pike	3	0	2	1	no limit
Walleye	0	3	1	0.5	no limit
<u>Chinook River</u>					
Walleye	2	3	1	0.5	no limit
<u>Crowley Reservoir</u>					
Northern pike	0	avoid	avoid	avoid	5
<u>Crowley Valley Reservoir</u>					
Northern pike	3	3	1.5	0.5	no limit
Walleye	2	avoid	avoid	avoid	0
<u>Crow (Steele) Lake</u>					
Northern pike	0	0	2.5	1.5	no limit
<u>Deer Lake</u>					
Northern pike	0	0	2	1	no limit
Walleye	2	avoid	avoid	avoid	5
<u>Endicott Lake</u>					
Walleye	3	0	2.5	1.5	no limit
<u>Eraser Lake</u>					
Northern pike	0	avoid	avoid	avoid	5
Walleye	0	avoid	avoid	avoid	0
<u>Etter Lake</u>					
Northern pike	2	3	1	0.5	no limit
Walleye	2	3	1	0.5	no limit
<u>Griffith Lake</u>					
Northern pike	3	3	1	0.5	no limit
Walleye	3	0	2	1	no limit
<u>Heide Lake</u>					
Northern pike	2	2	1	0.5	0
Walleye	2	avoid	avoid	avoid	5
<u>Isle Lake</u>					
Walleye	3	0	2.5	1.5	no limit
Northern pike	7	3	1	0.5	no limit
<u>Katowen Lake</u>					
Northern pike	0	3	1.5	0.5	no limit
Walleye	2	0	2	1	no limit
<u>Katie Lake</u>					
Walleye	0	3	2	1	no limit
<u>Kennedy Lake</u>					
Northern pike	3	3	1	0.5	no limit
Walleye	2	avoid	avoid	avoid	5

Water Body/Fish Species	Fish Size over lbs	Consumption Limit (servings per week)			
		Women	Children (5-11 yr)	Children (1-4 yr)	Adults
<u>Lac Beauport</u>					
Walleye	2	3	1	0.5	no limit
<u>Lac la Poudre</u>					
Northern pike	2	4	2	1	no limit
Walleye	2	avoid	avoid	avoid	6
<u>Lake Athabasca</u>					
Lake trout	6	5	2	1	no limit
Northern pike	6	5	2	1	no limit
Walleye	3	5	2	1	no limit
<u>Lesser Slave Lake</u>					
Northern pike	5	3	1	0.5	no limit
Walleye	3	4	2	1	no limit
<u>Little Bow Reservoir</u>					
Northern pike	3	2	1	0.5	6
<u>Lorne Lake</u>					
Northern pike	2	4	2	1	no limit
<u>Loun River</u>					
Walleye	2	4	2	1	no limit
<u>Marie Lake</u>					
Northern pike	4	5	2	1	no limit
<u>McGregor Lake</u>					
Walleye	4	avoid	avoid	avoid	5
<u>McMillan Lake</u>					
Northern pike	4	3	1	0.5	no limit
<u>Moose Lake</u>					
Northern pike	5	3	1	0.5	no limit
Walleye	4	2	1	0.5	no limit
<u>Murf's Watrous Lake</u>					
Northern pike	9	4	2	1	no limit
Walleye	4	4	2	1	no limit
<u>Pine Lake</u>					
Northern pike	3	5	2	1	no limit
Walleye	2	4	2	1	no limit
<u>Pinehurst Lake</u>					
Northern pike	3	avoid	avoid	avoid	4
Walleye	4	avoid	avoid	avoid	5
<u>Pithers Lake</u>					
Lake trout	10	avoid	avoid	avoid	4
<u>Richardson Lake¹</u>					
Northern pike	9	4	2	1	no limit
Walleye	3	5	2	1	no limit
<u>Rolling Hills Reservoir</u>					
Northern pike	8	avoid	avoid	avoid	3
Walleye	6	avoid	avoid	avoid	3
<u>Sylvan Lake</u>					
Walleye	1	6	2.5	1.5	no limit
<u>Touchwood Lake</u>					
Northern pike	10	avoid	avoid	avoid	4
Walleye	5	avoid	avoid	avoid	3
<u>Whitman Lake</u>					
Northern pike	7	3	1	0.5	no limit

Water Body/Fish Species	Fish Sized over lbs	Consumption Limit (servings per week)			
		Women	Children (5-11 yr)	Children (1-4 yr)	Adults
<u>Whitfish Lake</u>					
Lake whitefish	8	avoid	avoid	avoid	5
Northern pike	4	avoid	avoid	avoid	5
Walleye	4	avoid	avoid	avoid	4
<u>Wizard Lake</u>					
Northern pike	2	6	2.5	1.5	no limit
<u>Wolf Lake</u>					
Northern pike	3	4	2	1	no limit
Walleye	2	avoid	avoid	avoid	5

*1 lb = 454 grams. **1 serving = 75 grams, 1/2 cup, 2.5 ounces, or a piece of cooked fish that fits into the palm of your hand. *** "Women" refers women at reproductive age (15-49 yr) and pregnant women. Adult* includes adults and child over 12 yr. * calculations based on data from Environment Canada

5. References

- AHW (Alberta Health and Wellness). Swan Hills Special Waste Treatment Centre - Human Health Impact Assessment. Edmonton, Alberta, Canada, (1997).
- AHW (Alberta Health and Wellness). Wabamun and Area Community Exposure and Health Effects Assessment Program. Edmonton, Alberta, Canada: Alberta Health and Wellness, (2006).
- AHW (Alberta Health and Wellness). Human Health Risk Assessment - Mercury in Fish in Rivers and Lakes, Southern Alberta. Edmonton, Alberta: Alberta Health and Wellness, (2009a).
- AHW (Alberta Health and Wellness). Human Health Risk Assessment - Mercury in Fish in Southern Alberta Pine Coulee and Twin Valley Water Management Project. Edmonton, Alberta: Alberta Health and Wellness, (2009b).
- AHW (Alberta Health and Wellness). Human Health Risk Assessment - Mercury in Fish in Central Alberta Lac la Nonne and Lac Ste. Anne. Edmonton, Alberta: Alberta Health and Wellness, (2009c).
- AHW (Alberta Health and Wellness). Human Health Risk Assessment - Mercury in Fish - RAMP. Edmonton, Alberta: Alberta Health and Wellness, (2009d).
- Bhavsar, SP, Gewurtz, SB, McGoldrick, DJ, Keir, MJ, Backus, SM. Changes in mercury levels in Great Lakes fish between 1970s and 2007. *Environ. Sci. Technol.* 44: 3273-3279 (2010).
- Bodaly RA, St Louis VL, Paterson MJ, Fudge RJ, Hall BD, Rosenberg DM, Rudd JW. Bioaccumulation of mercury in the aquatic food chain in newly flooded areas. *Met Ions Biol Syst* 34:259-287 (1997).
- Bodaly RA, Jansen WA, Majewski AR, Fudge RJ, Strange NE, Derksen AJ, Green DJ. Postimpoundment time course of increased mercury concentrations in fish in hydroelectric reservoirs of northern Manitoba, Canada. *Arch Environ Contam Toxicol* 53:379-389 (2007).
- Bouzan C, Cohen JT, Connor WE, Kris-Etherton PM, Gray GM, Konig A, Lawrence RS, Savitz DA, Teutsch SM. A quantitative analysis of fish consumption and stroke risk. *Am J Prev Med* 29:347-352 (2005).
- Braune B, Muir D, DeMarch B, Gamberg M, Poole K, Currie R, Dodd M, Duschenko W, Eamer J, Elkin B, Evans M, Grundy S, Hebert C, Johnstone R, Kidd K, Koenig B, Lockhart L, Marshall H, Reimer K, Sanderson J, Shutt L. Spatial and temporal trends of contaminants in Canadian Arctic freshwater and terrestrial ecosystems: a review. *Sci Total Environ* 230:145-207 (1999).

Brinkmann, L, Rasmussen, JB. High levels of mercury in biota of a new Prairie irrigation reservoir with a simplified food web in Southern Alberta, Canada. *Hydrobiologia* 641: 11-21 (2010).

Cabana G, Tremblay A, Rasmussen JB. Pelagic food chain structure in Ontario Lakes: a determinant of mercury levels in Lake Trout. *Canadian Journal of Fisheries and Aquatic Sciences* 51:381-389 (1994).

Canadian Food Inspection Agency (CFIA). Draft Sablefish Mercury Report - Investigation of mercury in B.C. Sablefish sampled between October 2002 and November 2003. Unpublished document. (2003).

Chan HM, Egeland GM. Fish consumption, mercury exposure, and heart diseases. *Nutr Rev* 62:68-72 (2004).

Cohen JT, Bellinger DC, Connor WE, Shaywitz BA. A quantitative analysis of prenatal intake of n-3 polyunsaturated fatty acids and cognitive development. *Am J Prev Med* 29:366-374 (2005a).

Cohen JT, Bellinger DC, Shaywitz BA. A quantitative analysis of prenatal methyl mercury exposure and cognitive development. *Am J Prev Med* 29:353-365 (2005b).

Cohen JT, Bellinger DC, Connor WE, Kris-Etherton PM, Lawrence RS, Savitz DA, Shaywitz BA, Teutsch SM, Gray GM. A quantitative risk-benefit analysis of changes in population fish consumption. *Am J Prev Med* 29:325-334 (2005c).

Debes F, Budtz-Jorgensen E, Weihe P, White RF, Grandjean P. Impact of prenatal methylmercury exposure on neurobehavioral function at age 14 years. *Neurotoxicol Teratol* 28:536-547 (2006).

Domingo JL. Omega-3 fatty acids and the benefits of fish consumption: is all that glitters gold? *Environ Int* 33:993-998 (2007).

Dovydaitis T. Fish consumption during pregnancy: an overview of the risks and benefits. *J Midwifery Women Health* 53:325-330 (2008).

Dunstan JA, Simmer K, Dixon G, Prescott SL. Cognitive assessment of children at age 2(1/2) years after maternal fish oil supplementation in pregnancy: a randomised controlled trial. *Arch Dis Child Fetal Neonatal* Ed 93:F45-F50 (2008).

EHSUA (Environmental Health Sciences Program UoA. Development of A Fish Consumption Advisory Risk Communication/Public Participation Protocol for Alberta: A Case Study in Athabasca River and Tributaries. unpublished, internal report, 2000.

Evans, MS, Talbot, A. Investigation of mercury concentrations in walley and other fish in the Athabasca River ecosystem with increasing oil sands developments. J. Environ. Monitor. 14: 1989-2003.

Evans MS, Muir D, Lockhart WL, Stern G, Ryan M, Roach P. Persistent organic pollutants and metals in the freshwater biota of the Canadian Subarctic and Arctic: an overview. Sci Total Environ 351-352:94-147 (2005a).

Evans MS, Lockhart WL, Doetzel L, Low G, Muir D, Kidd K, Stephens G, Delaronde J. Elevated mercury concentrations in fish in lakes in the Mackenzie River Basin: the role of physical, chemical, and biological factors. Sci Total Environ 351-352:479-500 (2005b).

Feeley MM, Lo M-T. Risk Assessment for Mercury in Health Canada - Development of the Provisional Tolerable Daily Intake (pTDI) Value. In: Proceedings of the Conference on Mercury in Eastern Canada and the Northeast States, 21-23 September, 1998 (Pilgrim W, Burgess N, Giguère M-F, eds).(1998).

Feeley MM. Health Canada Mercury Guidelines. Personal communication.

Garcia E, Carignan R. Mercury concentrations in fish from forest harvesting and fire-impacted Canadian Boreal lakes compared using stable isotopes of nitrogen. Environ Toxicol Chem 24:685-693 (2005).

Gerstenberger SL, Dellinger JA. PCBs, mercury, and organochlorine concentrations in lake trout, walleye, and whitefish from selected tribal fisheries in the Upper Great Lakes region. Environ Toxicol 17:513-519 (2002).

Hallgren CG, Hallmans G, Jansson JH, Marklund SL, Huhtasaari F, Schutz A, Stromberg U, Vessby B, Skerfving S. Markers of high fish intake are associated with decreased risk of a first myocardial infarction. Br J Nutr 86:397-404 (2001).

Health Canada. Methylmercury in Canada. Exposure of Indian and Inuit Residents to Methylmercury in the Canadian Environment. Ottawa:Ministry of National Health and Welfare, Medical Services Branch, (1979).

Health Canada. Human Health Risk Assessment of Mercury in Fish and Health Benefits of Fish Consumption. Ottawa, Canada:Health Canada, (2007).

http://www.hc-sc.gc.ca/fn-an/pubs/mercur/merc_fish_poisson-eng.php

Health Canada. Lesser Slave Lake Health Study. Unpublished, Medical Service Branch, Health Canada, (1999).

Helland IB, Smith L, Saarem K, Saugstad OD, Drevon CA. Maternal supplementation with very-long-chain n-3 fatty acids during pregnancy and lactation augments children's IQ at 4 years of age. Pediatrics 111:e39-e44 (2003).

Jewett SC, Zhang X, Naidu AS, Kelley JJ, Dasher D, Duffy LK. Comparison of mercury and methylmercury in Northern pike and Arctic grayling from western Alaska rivers. *Chemosphere* 50:383-392 (2003).

Kamman NC, Burgess NM, Driscoll CT, Simonin HA, Goodale W, Linehan J, Estabrook R, Hutcheson M, Major A, Scheuhammer AM, Scruton DA. Mercury in freshwater fish of northeast North America—a geographic perspective based on fish tissue monitoring databases. *Ecotoxicology* 14:163-180 (2005).

Kerper LE, Ballatori N, Clarkson TW. Methylmercury transport across the blood-brain barrier by an amino acid carrier. *Am J Physiol* 262:R761-R765 (1992).

Kinghorn A, Solomon P, Chan HM. Temporal and spatial trends of mercury in fish collected in the English-Wabigoon river system in Ontario, Canada. *Sci Total Environ* 372:615-623 (2007).

Kirk JL, Muir DC, Gleason A, Wang X, Lawson G, Frank RA, Lehnher I, Wrona F. Atmospheric deposition of mercury and methylmercury to landscapes and waterbodies of the Athabasca oil sands region. *Environ. Sci. Technol.* 48(13):7374-83 (2014)

Konig A, Bouzan C, Cohen JT, Connor WE, Kris-Etherton PM, Gray GM, Lawrence RS, Savitz DA, Teutsch SM. A quantitative analysis of fish consumption and coronary heart disease mortality. *Am J Prev Med* 29:335-346 (2005).

Kris-Etherton PM, Harris WS, Appel LJ. Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease. *Arterioscler Thromb Vasc Biol* 23:e20-e30 (2003).

Levenson CW, Axelrad DM. Too much of a good thing? Update on fish consumption and mercury exposure. *Nutr Rev* 64:139-145 (2006).

Lockhart WL, Stern GA, Low G, Hendzel M, Boila G, Roach P, Evans MS, Billeck BN, Delaronde J, Friesen S, Kidd K, Atkins S, Muir DC, Stoddart M, Stephens G, Stephenson S, Harbicht S, Snowshoe N, Grey B, Thompson S, DeGraff N. A history of total mercury in edible muscle of fish from lakes in northern Canada. *Sci Total Environ* 351-352:427-463 (2005).

Mahaffey KR, Clickner RP, Jeffries RA. Methylmercury and omega-3 fatty acids: co-occurrence of dietary sources with emphasis on fish and shellfish. *Environ Res* 107:20-29 (2008).

McDowell, MA, Dillon, CF, Osterloh, J, P. Bolger, M, Pellizzari, E, Fernando, R, Montes de Oca, R, Schober, SE, Sinks, T, Jones, RL, Mahaffey KR. Hair mercury levels in U.S. children and women of childbearing age: Reference range data from NHANES 1999-2000. *Environ Health Persp* 112: 1165-1171.

Mergler D, Anderson HA, Chan LH, Mahaffey KR, Murray M, Sakamoto M, Stern AH. Methylmercury exposure and health effects in humans: a worldwide concern. *Ambio* 36:3-11 (2007).

Morgan JN, Berry MR, Graves RL. Effects of commonly used cooking practices on total mercury concentration in fish and their impact on exposure assessments. *J Expo Anal Environ Epidemiol* 7:119-133 (1997).

Mozaffarian D, Shi P, Morris JS, Grandjean P, Siscovick DS, Spiegelman D, Willett WC, Rimm EB, Curhan GC, Forman JP. Mercury exposure and risk of hypertension in US men and women in 2 prospective cohorts. *Hypertension*, 60(3): 645-652 (2012).

Mozaffarian D, Rimm EB. Fish intake, contaminants, and human health: evaluating the risks and the benefits. *JAMA* 296:1885-1899 (2006).

Munn MB, Short TM. Spatial heterogeneity of mercury bioaccumulation by walleye in Franklin D. Roosevelt Lake and the Upper Columbia River, Washington. *Trans Am Fish Soc* 126:477-487 (1997).

Neff MR, Bhavsar SP, Arhonditsis GB, Fletcher R, Jackson DA. Long term changes in fish mercury levels in the historically impacted English – Wabigoon River system (Canada). *J Environ. Monit.* 14: 2327-2337 (2012).

Neumann RM, Ward SM. Bioaccumulation and biomagnification of mercury in two warmwater fish communities. *J Freshwater Ecol* 14:487-494 (1999).

Oken E, Kleinman KP, Berland WE, Simon SR, Rich-Edwards JW, Gilman MW. Decline in fish consumption among pregnant women after a national mercury advisory. *Obstet Gynecol* 102:346-351 (2003).

Oken E, Bellinger DC. Fish consumption, methylmercury and child neurodevelopment. *Curr Opin Pediatr* 20:178-183 (2008).

Oken E, Radesky JS, Wright RO, Bellinger DC, Amarasingwardena CJ, Kleinman KP, Hu H, Gilman MW. Maternal fish intake during pregnancy, blood mercury levels, and child cognition at age 3 years in a US cohort. *Am J Epidemiol* 167:1171-1181 (2008).

Schoeman K, Bernd JR, Hill J, Nash K, Koren G. Defining a lowest observable adverse effect hair concentration of mercury for neurodevelopmental effects of prenatal methylmercury exposure through maternal fish consumption: A systematic review. *Ther Drug Monitor* 31: 670-682 (2009).

Selch TM, Hoagstrom CW, Weimer EJ, Duehr JP, Chipps SR. Influence of fluctuating water levels on mercury concentrations in adult walleye. *Bull Environ Contam Toxicol* 79:36-40 (2007).

Stern AH. A review of the studies of the cardiovascular health effects of methylmercury with consideration of their suitability for risk assessment. *Environ Res* 98:133-142 (2005).

Suzuki SC, Sorensen JA, Peddicord RK, Clifford PA, Dillon TM. Trophic transfer and biomagnification potential of contaminants in aquatic ecosystems. *Rev Environ Contam Toxicol* 136:21-89 (1994).

Uday R, Huffman DR, Mena P, Llanos A, Birch EE. Term infant studies of DHA and ARA supplementation on neurodevelopment: results of randomized controlled trials. *J Pediatr* 143:S17-S25 (2003).

USEPA (United States Environmental Protection Agency). Trace Elements in Water, Solids, And Biosolids By Inductively Coupled Plasma-Atomic Emission Spectrometry EPA-821-R-01-010. Washington DC Office of Sciences and Technology, USEPA, (2001).

USEPA (United States Environmental Protection Agency). Guidance for Assessing Chemical Contaminant Data for Use in Fish Advisories & Risk Assessment and Fish Consumption Limits. EPA 823-B-00-008. Washington, DC Office of Science and Technology, Office of Water, USEPA, (2000).

USEPA (United States Environmental Protection Agency). Mercury in Solids and Solutions by Thermal Decomposition, Amalgamation, and Atomic Absorption Spectrophotometry. Method 1631. Washington, DC Office of Science and Technology, USEPA, (2007).

Verbeke W, Verhoyden F, Fremer LJ, Sioen I, De Henauw S, Van Camp J. Communicating Risks and Benefits from Fish Consumption: Impact on Belgian Consumers' Perception and Intention to Eat Fish. *Risk Anal* 28:951-957 (2008).

Vinton R, Brouard D, Demers C, Leclerc R, Lesard M, Scheraga R. Mercury evolution (1978-1988) in fishes of the La Grande Hydroelectric complex, Quebec, Canada. *Water, Air, and Soil Pollution* 56:405-417 (1991).

Vitonen JR, Laakkonen JA, Mursu J, Vuolteenaho S, Tuomariemi T-P. Serum long-chain n-3 polyunsaturated fatty acids, mercury, and risk of sudden cardiac death in men: A prospective population-based study. *PLoS ONE* 7(7): e41046 (2012).

Vitonen JR, Vuolteenaho S, Rissanen TH, Mursu J, Tuomariemi T-P, Kujala M, Vauhonen V-P, Seppänen R, Laakkonen JA, Salonen JT. Mercury, fish oils, and risk of acute coronary events and cardiovascular disease, coronary heart disease, and all-cause mortality in men in eastern Finland. *Atherosclerosis, Thrombosis, and Vascular Biology* 25: 228-233 (2005).

WHO (World Health Organization). Methylmercury. Geneva:World Health Organization, (1990).

Yang, D.-W., Yu, X., Chen, Y.-H., Sebaste, K. Inverse relationships between selenium and mercury in tissues of young walleye (*Stizostedion vitreum*) from Canadian inland lakes. *Sci. Tot. Environ.* 438: 1075–1082 (2012).

Zhang, L., Campbell, L.B., Johnson, T.B. Seasonal variation in mercury and food web biomagnification in Lake Ontario. *Environ. Pollut.* 151: 175–186 (2012).

Zhang, J., Smith, S., Amaro, E., Kozlowski, H. Fish consumption and mortality from all causes, ischemic heart disease, and stroke: an ecological study. *Proc. Natl. Acad. Sci.* 96: 5275–5279 (1999).

Supplement 6

Special Issue: Environmental Health



[illegible]

Appendix B

Guide for Use of Food Consumption Advisory Information

1. Where to find food consumption advisory information?

Fish consumption advisories

<http://www.alberta.ca/fishing-safety-procedures/fish-consumption-advisory.aspx>

Wild game meat consumption advisories

<http://www.alberta.ca/fishing-safety-procedures/wild-game-meat-consumption-advisory.aspx>

2. What are criteria for issuing food consumption advisories?

The criteria for issuing fish consumption advisories are

1. If mercury levels are higher than 0.5 mg/kg (commercial fishing guideline), the advisory would be "avoid consuming fish",
2. If mercury levels are between 0.2 - 0.5 mg/kg (Health Canada recommendation for subsistence consumers), the advisory would provide "consumption limits",
3. If mercury levels are less than 0.2 mg/kg, advisory would not be issued,
4. If the fish sample size is less than 5 per location, advisory would not be issued, and
5. If the lakes are used for commercial fishing, advisories would not be issued until consulting with Canadian Food Inspection agency.

The criteria for issuing wild game meat consumption advisories are based on the guidelines proposed by national or international regulatory agencies such as Health Canada and World Health Organization.

3. How to find fish species and size?

The fish species are indicated under the "Species" in the table. Meanwhile, please check fish weight under the "Fish Size" in the table. If fish weighs less than the weight indicated in the table, there is no advisory against eating this size of fish.

4. How to find meal size and frequency?

Please go to the "Consumption Limits", the meal size and frequency indicated as "servings per week". Check specific information on meal size and frequency for women, young children groups and adults.